

JC14 Rec'd PCT/PTO 01 OCT 2001

FORM PTO-1390 (REV. 11-2000)		U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE		ATTORNEY'S DOCKET NUMBER 0230-0169P	
TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 U.S.C. 371					
INTERNATIONAL APPLICATION NO. PCT/JP00/02080		INTERNATIONAL FILING DATE March 31, 2000		U.S. APPLICATION NO. (If known, see 37 CFR 1.52) 09/937905 NEW	
TITLE OF INVENTION NOVEL PROTEINS GENE ENCODING THE SAME AND METHOD OF UTILIZATION THEREOF		PRIORITY DATE CLAIMED April 1, 1999			
APPLICANT(S) FOR DO/EO/US SHA, Shigen; AOKI, Yoshihiko; NISHI, Yoshisuke					
Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:					
1. <input checked="" type="checkbox"/> This is a <b>FIRST</b> submission of items concerning a filing under 35 U.S.C. 371. 2. <input type="checkbox"/> This is a <b>SECOND</b> or <b>SUBSEQUENT</b> submission of items concerning a filing under 35 U.S.C. 371. 3. <input checked="" type="checkbox"/> This express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39 (1). 4. <input checked="" type="checkbox"/> The US has been elected by the expiration of 19 months from the priority date (Article 31). 5. <input checked="" type="checkbox"/> A copy of the International Application as filed (35 U.S.C. 371(c)(2)) a. <input type="checkbox"/> is transmitted herewith (required only if not transmitted by the International Bureau). b. <input checked="" type="checkbox"/> has been transmitted by the International Bureau. WO 00/60075 c. <input type="checkbox"/> is not required, as the application was filed in the United States Receiving Office (RO/US). 6. <input checked="" type="checkbox"/> An English language translation of the International Application as filed (35 U.S.C. 371(c)(2)). a. <input checked="" type="checkbox"/> is transmitted herewith. b. <input type="checkbox"/> has been previously submitted under 35 U.S.C. 154(d)(4) 7. <input checked="" type="checkbox"/> Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3)). a. <input type="checkbox"/> are transmitted herewith (required only if not transmitted by the International Bureau). b. <input type="checkbox"/> have been transmitted by the International Bureau. c. <input type="checkbox"/> have not been made; however, the time limit for making such amendments has NOT expired. d. <input checked="" type="checkbox"/> have not been made and will not be made. 8. <input type="checkbox"/> An English language translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)). 9. <input checked="" type="checkbox"/> An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)). 10. <input type="checkbox"/> An English language translation of the annexes of the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).					
Items 11. to 20. below concern document(s) or information included:					
11. <input checked="" type="checkbox"/> An Information Disclosure Statement under 37 CFR 1.97 and 1.98, Form PTO-1449(s), and International Search Report (PCT/ISA/210). 12. <input type="checkbox"/> An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included. 13. <input checked="" type="checkbox"/> A <b>FIRST</b> preliminary amendment. 14. <input type="checkbox"/> A <b>SECOND</b> or <b>SUBSEQUENT</b> preliminary amendment. 15. <input type="checkbox"/> A substitute specification. 16. <input type="checkbox"/> A change of power of attorney and/or address letter. 17. <input type="checkbox"/> A computer-readable form of the sequence listing in accordance with PCT Rule 13ter.2 and 35 U.S.C. 1.821-1.825. 18. <input type="checkbox"/> A second copy of the published international application under 35 U.S.C. 154(d)(4). 19. <input type="checkbox"/> A second copy of the English language translation of the international application under 35 U.S.C. 154(d)(4). 20. <input checked="" type="checkbox"/> Other items or information: 1.) Verification of Translation 2.) PCT Request 3.) International Preliminary Examination Report (PCT/IPEA/409) 4.) Six (6) pages of Sequence Listing 5.) Zero (0) sheets of Formal Drawings					

JC05 Rec'd PCT/PTO 01 OCT 2001

U.S. APPLICATION NO (if known, see 37 CFR 1.5) <div style="font-size: 1.5em; font-weight: bold;">09/937905</div>		INTERNATIONAL APPLICATION NO PCT/JP00/02080		ATTORNEY'S DOCKET NUMBER 0230-0169P	
---	--	--	--	--	--

21. <input checked="" type="checkbox"/> The following fees are submitted: <b>BASIC NATIONAL FEE (37 CFR 1.492(a)(1)-(5):</b> Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO and International Search Report not prepared by the EPO or JPO. .... <b>\$1,040.00</b>  International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International Search Report prepared by the EPO or JPO ..... <b>\$890.00</b>  International preliminary examination fee (37 CFR 1.482) not paid to USPTO but international search fee (37 CFR 1.445(a)(2)) paid to USPTO. .... <b>\$740.00</b>  International preliminary examination fee (37 CFR 1.482) paid to USPTO but all claims did not satisfy provisions of PCT Article 33(1)-(4) ..... <b>\$710.00</b>  International preliminary examination fee (37 CFR 1.482) paid to USPTO and all claims satisfied provisions of PCT Article 33(1)-(4). .... <b>\$100.00</b> <b>ENTER APPROPRIATE BASIC FEE AMOUNT =</b>	<b>CALCULATIONS      PTO USE ONLY</b>							
Surcharge of <b>\$130.00</b> for furnishing the oath or declaration later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(e)).	<table style="width:100%; border-collapse: collapse;"> <tr> <td style="width:10%; text-align: center;">\$</td> <td style="width:40%; text-align: right;">890.00</td> <td style="width:50%;"></td> </tr> <tr> <td style="text-align: center;">\$</td> <td style="text-align: right;">0</td> <td></td> </tr> </table>		\$	890.00		\$	0	
\$	890.00							
\$	0							

CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE		
Total Claims	50 - 20 =	30	X \$18.00	\$	540.00
Independent Claims	10 - 3 =	7	X \$84.00	\$	588.00
MULTIPLE DEPENDENT CLAIM(S) (if applicable) Yes			+ \$280.00	\$	280.00
<b>TOTAL OF ABOVE CALCULATIONS =</b>				\$	2298.00

<input type="checkbox"/> Applicant claims small entity status. See 37 CFR 1.27. The fees indicated above are reduced by 1/2.	<table style="width:100%; border-collapse: collapse;"> <tr> <td style="width:10%; text-align: center;">\$</td> <td style="width:40%; text-align: right;">0</td> <td style="width:50%;"></td> </tr> </table>		\$	0	
\$	0				
<b>SUBTOTAL =</b>			\$	2298.00	

Processing fee of <b>\$130.00</b> for furnishing the English translation later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(f)).	<table style="width:100%; border-collapse: collapse;"> <tr> <td style="width:10%; text-align: center;">\$</td> <td style="width:40%; text-align: right;">0</td> <td style="width:50%;"></td> </tr> </table>		\$	0	
\$	0				
<b>TOTAL NATIONAL FEE =</b>			\$	2298.00	

Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). <b>\$40.00</b> per property +	<table style="width:100%; border-collapse: collapse;"> <tr> <td style="width:10%; text-align: center;">\$</td> <td style="width:40%; text-align: right;">0</td> <td style="width:50%;"></td> </tr> </table>		\$	0	
\$	0				
<b>TOTAL FEES ENCLOSED =</b>			\$	2298.00	

	Amount to be:	
	refunded	
	charged	

a. ☒ A check in the amount of \$ **2298.00** to cover the above fees is enclosed.

b. ☐ Please charge my Deposit Account. No. \_\_\_\_\_ in the amount of \$ \_\_\_\_\_ to cover the above fees.  
 A duplicate copy of this sheet is enclosed.

c. ☒ The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. 02-2448.

**NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.**

Send all correspondence to:  
**Birch, Stewart, Kolasch & Birch, LLP** or Customer No. 2292  
**P.O. Box 747**  
**Falls Church, VA 22040-0747**  
**(703) 205-8000**

**Date: October 1, 2001**

By *Gerard M. Murphy, Jr.* #28,977  
 Gerard M. Murphy, Jr., #28,977



09937905 09/1787970  
Rec'd PCT/PTO 14 JAN 2002  
BOX SEQUENCE  
PATENT  
230-169P

IN THE U.S. PATENT AND TRADEMARK OFFICE

Applicant: Shiken SHA et al. Conf.: 5513  
Appl. No.: 09/937,905 Group: 1645  
Filed: October 1, 2001 Examiner: TO BE ASSIGNED  
For: NOVEL PROTEINS, GENE ENCODING THE SAME AND  
METHOD OF UTILIZATION THEREOF

AMENDMENT

Assistant Commissioner for Patents  
Washington, DC 20231

January 14, 2002

Sir:

In response to the Notification of Missing Requirements Under 35 U.S.C. 371 in the United States Designated/Elected Office (DO/EO/US) mailed November 14, 2001, the following amendments and remarks are respectfully submitted in connection with the above-identified application.

In the Specification:

Please replace the paragraph beginning on page 3, line 17, with the following rewritten paragraph:

In other words, the present invention provides a gene encoding: (a) a protein having the amino acid sequence listed as SEQ ID NO:2 of the Sequence Listing; (b) a protein having the amino acid sequence listed as SEQ ID NO:2 of the Sequence Listing with one or more amino acid deletions, substitutions, additions or insertions and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor; or (c) a protein having at least 50% homology with the amino acid sequence listed as SEQ ID NO:2 and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor.

Application No. 09/937,905

Please replace the paragraph beginning on page 4, line 1, with the following rewritten paragraph:

The invention further provides a gene encoding: (a) a protein having the amino acid sequence listed as SEQ ID NO:4 of the Sequence Listing; (b) a protein having the amino acid sequence listed as SEQ ID NO:4 of the Sequence Listing with one or more amino acid deletions, substitutions, additions or insertions and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor; or (c) a protein having at least 50% homology with the amino acid sequence listed as SEQ ID NO:4 and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor.

Please replace the paragraph beginning on page 4, line 13, with the following rewritten paragraph:

The invention further provides a gene having: (a) the nucleotide sequence listed as SEQ ID NO:1 of the Sequence Listing; (b) a nucleotide sequence which is the nucleotide sequence listed as SEQ ID NO:1 of the Sequence Listing with one or more nucleotide deletions, substitutions, additions or insertions and which encodes a protein that can bind to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor; or (c) a nucleotide sequence which hybridizes with DNA having the nucleotide sequence listed as SEQ ID NO:1 of the Sequence Listing under stringent conditions and encodes a protein that can bind to an antibody or its fragments that are active to induce granulocyte colony-stimulating factor.

Please replace the paragraph beginning on page 4, line 26, with the following rewritten paragraph:

The invention further provides a gene having: (a) the nucleotide sequence listed as SEQ ID NO:3 of the Sequence Listing; (b) a nucleotide sequence which is the nucleotide sequence listed as SEQ ID NO:3 of the Sequence Listing with one or more nucleotide deletions, substitutions, additions or insertions and which encodes a protein that can bind to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor; or (c) a nucleotide sequence which hybridizes with DNA having the nucleotide sequence listed as SEQ ID NO:3 of the Sequence Listing under stringent conditions and encodes a protein that can bind to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor.

Application No. 09/937,905

Please replace the paragraph beginning on page 5, line 17, with the following rewritten paragraph:

The invention further provides a DNA fragment containing: (1) the nucleotide sequence from position 519 to position 736, the nucleotide sequence from position 666 to position 689, the nucleotide sequence from position 381 to position 403 or the nucleotide sequence from position 709 to position 727 of the nucleotide sequence listed as SEQ ID NO:1 of the Sequence Listing; (2) a nucleotide sequence which is any of the nucleotide sequences of (1) above with one or more nucleotide deletions, substitutions, additions or insertions; or (3) a nucleotide sequence which has at least 80% homology with any of the nucleotide sequences of (1) above.

Please replace the paragraph beginning on page 6, line 1, with the following rewritten paragraph:

The invention further provides a gene containing: (1) the nucleotide sequence from position 519 to position 736, the nucleotide sequence from position 666 to position 689, the nucleotide sequence from position 381 to position 403 or the nucleotide sequence from position 709 to position 727 of the nucleotide sequence listed as SEQ ID NO:1 of the Sequence Listing; (2) a nucleotide sequence which is any of the nucleotide sequences of (1) above with one or more nucleotide deletions, substitutions, additions or insertions; or (3) a nucleotide sequence which has at least 80% homology with any of the nucleotide sequences of (1) above; and encoding a protein that can bind to an antibody or its fragments that are active to induce granulocyte colony-stimulating factor.

Please replace the paragraph beginning on page 6, line 15, with the following rewritten paragraph:

The invention further provides any of the following proteins: (a) a protein having the amino acid sequence listed as SEQ ID NO:2 of the Sequence Listing; (b) a protein having the amino acid sequence listed as SEQ ID NO:2 of the Sequence Listing with one or more amino acid deletions, substitutions, additions or insertions and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor; (c) a protein having at least 50% homology with the amino acid sequence listed as SEQ ID NO:2 and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor; or (d) a protein that is encoded by DNA which hybridizes with DNA having the nucleotide sequence listed as SEQ ID NO:1 of the Sequence

Application No. 09/937,905

Listing under stringent conditions and that binds to an antibody or its fragments that are active to induce granulocyte colony-stimulating factor.

Please replace the paragraph beginning on page 7, line 4, with the following rewritten paragraph:

The invention further provides any of the following proteins: (a) a protein having the amino acid sequence listed as SEQ ID NO:4 of the Sequence Listing; (b) a protein having the amino acid sequence listed as SEQ ID NO:4 of the Sequence Listing with one or more amino acid deletions, substitutions, additions or insertions and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor; (c) a protein having at least 50% homology with the amino acid sequence listed as SEQ ID NO:4 and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor; or (d) a protein that is encoded by DNA which hybridizes with DNA having the nucleotide sequence listed as SEQ ID NO:3 of the Sequence Listing under stringent conditions and that binds to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor.

Please replace the paragraph beginning on page 7, line 28, with the following rewritten paragraph:

The invention further provides a protein comprising any of the followings: (1) the amino acid sequence from residues 1 to 91, the amino acid sequence from residues 50 to 146, the amino acid sequence from residues 1 to 78, the amino acid sequence from residues 200 to 241, the amino acid sequence from residues 172 to 241, the amino acid sequence from residues 103 to 150 or the amino acid sequence from residues 169 to 241 of the amino acid sequence listed as SEQ ID NO:2 of the Sequence Listing; (2) an amino acid sequence which is any of the amino acid sequences of (1) above with one or more amino acid deletions, substitutions, additions or insertions; or (3) an amino acid sequence having at least 70% homology with any of the amino acid sequences of (1) above.

Please replace the paragraph beginning on page 8, line 14, with the following rewritten paragraph:

The invention further provides a protein comprising any of the followings: (1) the amino acid sequence from residues 1 to 91, the amino acid sequence from residues 50 to 146, the amino acid

Application No. 09/937,905

sequence from residues 1 to 78, the amino acid sequence from residues 200 to 241, the amino acid sequence from residues 172 to 241, the amino acid sequence from residues 103 to 150 or the amino acid sequence from residues 169 to 241 of the amino acid sequence listed as SEQ ID NO:2 of the Sequence Listing; (2) an amino acid sequence which is any of the amino acid sequences of (1) above with one or more amino acid deletions, substitutions, additions or insertions; or (3) an amino acid sequence having at least 70% homology with any of the amino acid sequences of (1) above, and also binding to an antibody or its fragments that are active to induce granulocyte colony-stimulating factor.

Please replace the paragraph beginning on page 17, line 5, with the following rewritten paragraph:

The present invention provides a gene which encodes the protein having the amino acid sequence listed as SEQ ID NO:1 of the Sequence Listing or a protein which is homologous thereto. The invention also provides a gene having the nucleotide sequence listed as SEQ ID NO:1 of the Sequence Listing or a nucleotide sequence which is homologous thereto.

Please replace the paragraph beginning on page 17, line 16, with the following rewritten paragraph:

The gene of the invention will typically have the nucleotide sequence listed as SEQ ID NO:1 of the Sequence Listing, but this is only the sequence of a clone (MMR19) obtained in the following examples to represent an embodiment of the invention. Those skilled in the art are well aware that natural genes have a small number of variations depending on the breeding conditions of the biological species that produce it, and on the ecosystem, or on the presence of highly similar isozymes. Consequently, the "gene" of the invention is not limited to the gene having the nucleotide sequence listed as SEQ ID NO:1 of the Sequence Listing but also includes all genes encoding proteins having the characters described in the present specification.

Please replace the paragraph beginning on page 19, line 1, with the following rewritten paragraph:

A homologous gene cloned by hybridization in this manner has at least 70%, preferably at least 80%, more preferably at least 90%, even more preferably at least 95% and most preferably at

least 98% homology with respect to the nucleotide sequence listed as SEQ ID NO:1 of the Sequence Listing.

Please replace the paragraph beginning on page 19, line 9, with the following rewritten paragraph:

The present invention provides the protein having the amino acid sequence listed as SEQ ID NO:1 of the Sequence Listing or a protein homologous thereto.

Please replace the paragraph beginning on page 19, line 12, with the following rewritten paragraph:

The protein having the amino acid sequence listed as SEQ ID NO:1 of the Sequence Listing according to the invention may be obtained by incorporating the gene encoding therefor into an appropriate expression vector, transforming this vectors to an appropriate host and expressing the recombinant protein. However, the source and preparation method are not restricted so long as the protein of the invention has the characters described in the present specification, and it may be a naturally produced protein, a protein expressed from recombinant DNA by a genetic engineering method or a protein chemically synthesized.

Please replace the paragraph beginning on page 19, line 24, with the following rewritten paragraph:

The protein of the invention will typically have the sequence of 241 amino acids listed as SEQ ID NO:1 of the Sequence Listing. However, those skilled in the art are well aware that natural proteins include variations of one or more amino acids due to gene variation depending on the breeding conditions of biological species that produce it, on the ecosystem, the presence of highly similar isozymes. The term "amino acid variation" as used here means one or more amino acid substitutions, deletions, insertions and/or additions. The "protein" of the invention has the amino acid sequence listed as SEQ ID NO:1 based on deduction from the nucleotide sequence of the cloned gene, but it is not limited only to proteins with that sequence and is intended to include all homologous proteins that have the characteristics described in the present specification. The homology is at least 50%, preferably at least 60%, more preferably at least 70%, more preferably at



least 80%, even more preferably at least 90%, yet more preferably at least 95% and most preferably at least 98%.

Please replace the paragraph beginning on page 21, line 16, with the following rewritten paragraph:

The total RNA is extracted from a human macrophage cell line (THP-1, U937, HL-60) by guanidium thiocyanate-phenol-chloroform single-step extraction (Laboratory Manuals of Genetic Engineering, 3rd Edition, pp.83-84, 1996), and purified using an oligo(dT) cellulose column, to obtain poly(A)<sup>+</sup> RNA. Reverse transcriptase (MMLV-RTase) and DNA polymerase are used to synthesize double-stranded cDNA. The double-stranded cDNA is used to construct a cDNA library using a  $\lambda$ ZAPII phage vector by the method of Gubler-Hoffmann (Gubler, U. and Hoffmann, B.J.: Gene, 25:263-269, 1983). A probe is then prepared by amplifying a DNA sequence using a primer DNA that can amplify a sequence in the region of the nucleotide sequence (SEQ ID NO:1) of the mouse cDNA (MMR19 clone) disclosed in the present specification, having high homology with the human sequence (for example, the region from position 172 to position 241 of SEQ ID NO:1 which has been found to have 91% homology with the human sequence) and the template DNA from the human macrophage cell cDNA library. Or region (for example, the region from position 172 to position 241 of SEQ ID NO:1) is used directly as the probe for screening the cDNA encoding the entire length of the target protein from the human macrophage cell cDNA library. The cDNA nucleotide sequence is analyzed by the Primer Walking method. The cDNA confirmed to encode the entire length of the target protein is introduced into a baculovirus to express a protein, which can be purified with an affinity column to obtain the human-type homologous protein.

Please replace the paragraph beginning on page 22, line 16, with the following rewritten paragraph:

As explained above, the present invention relates to the gene or protein having the nucleotide sequence listed as SEQ ID NO:1 or the amino acid sequence listed as SEQ ID NO:2, and to genes and proteins which are homologous thereto. As a result of a search to determine whether or not sequences homologous to the nucleotide sequence listed as SEQ ID NO:1 and the amino acid sequence listed as SEQ ID NO:2 provided by the invention are present in other organisms, it was confirmed that human ESTs (expressed sequence tags) include sequences having high homology

Application No. 09/937,905

with the gene of the invention (see Example 3 below). It will therefore be readily appreciated that the human-derived homologous gene can also be isolated by screening of a human-derived gene library (cDNA library, etc.) using the human-derived ESTs having high homology with the nucleotide sequence of the invention as the probe.

Please replace the paragraph beginning on page 23, line 4, with the following rewritten paragraph:

As described above, a database search revealed that portions (i.e. DNA fragments) of the nucleotide sequence listed as SEQ ID NO:1 according to the invention are conserved, having high homology with the human sequence. Such DNA fragments are useful as probes for screening of the human-derived homologous gene, and therefore constitute one aspect of the present invention. The DNA fragments include DNA fragments containing the nucleotide sequence from position 519 to position 736, the nucleotide sequence from position 666 to position 689, the nucleotide sequence from position 381 to position 403 or the nucleotide sequence from position 709 to position 727 of the nucleotide sequence listed as SEQ ID NO:1 of the Sequence Listing, while DNA fragments containing a nucleotide sequence which is any of these nucleotide sequences with one or more nucleotide deletions, substitutions, additions or insertions or a nucleotide sequence which has at least 80%, preferably at least 85%, more preferably at least 90%, even more preferably at least 95% and most preferably at least 98% homology with any of these nucleotide sequences, are also within the scope of the invention.

Please replace the paragraph beginning on page 23, line 25, with the following rewritten paragraph:

A database search also revealed that a portion of the amino acid sequence listed as SEQ ID NO:2 according to the invention is conserved with high homology in the human sequence. Protein fragments comprising portions of the protein of the invention are also useful as reagents for analysis and isolation of antibodies with G-CSF inducing activity, as is the protein of the invention, and also have potential utility as a drug like the protein of the invention, and thus constitute an aspect of the invention.

Application No. 09/937,905

Please replace the paragraph beginning on page 24, line 6, with the following rewritten paragraph:

The mentioned proteins include the amino acid sequences from residues 1 to 91, 50 to 146, 1 to 78, 200 to 241, 172 to 241, 103 to 150, and 169 to 241 of the amino acid sequence listed as SEQ ID NO:2 of the Sequence Listing, while proteins containing an amino acid sequence which is any of these amino acid sequences with one or more amino acid deletions, substitutions, additions or insertions or an amino acid sequence having at least 70%, preferably at least 80%, more preferably at least 85%, even more preferably at least 90%, even more preferably at least 95% and most preferably at least 98% homology with any of these amino acid sequences, are also within the scope of the invention.

Please replace the paragraph beginning on page 24, line 19, with the following rewritten paragraph:

The present inventors have determined the nucleotide sequence of the human-type antigen gene by a method similar to the one described above (see Example 4 below). Consequently, the present invention provides a gene having the nucleotide sequence listed as SEQ ID NO:3 of the Sequence Listing or having a nucleotide sequence which is homologous thereto. The invention also provides a protein having the amino acid sequence listed as SEQ ID NO:4 of the Sequence Listing or a protein which is homologous thereto. Here, "homologous" means that the scope of the invention is not limited to the gene having the nucleotide sequence listed as SEQ ID NO:3 or the protein having the amino acid sequence listed as SEQ ID NO:4, as was explained in the part of "Gene of the invention" and "Protein of the invention" described above.

Please replace the paragraph beginning on page 43, line 19, with the following rewritten paragraph:

The protein of the invention used for screening is preferably (a) a protein having the amino acid sequence listed as SEQ ID NO:4 of the Sequence Listing; (b) a protein having the amino acid sequence listed as SEQ ID NO:4 of the Sequence Listing with one or more amino acid deletions, substitutions, additions or insertions and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor; (c) a protein having at least 50% (preferably at least 60%, more preferably at least 70%, even more preferably at least 80%, even more preferably at least

Application No. 09/937,905

90%, especially preferably at least 94% and most preferably at least 98%) homology with the amino acid sequence listed as SEQ ID NO:4 and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor; or (d) a protein that is encoded by DNA which hybridizes with DNA having the nucleotide sequence listed as SEQ ID NO:3 of the Sequence Listing under stringent conditions and that binds to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor.

Please replace the paragraph beginning on page 55, line 9, with the following rewritten paragraph:

Based on the results of the nucleotide sequence analyses, a clone MMR19 was found to have the 840 bp nucleotide sequence of the full length cDNA which included the open reading frame of the protein. The nucleotide sequence of the clone MMR19 is listed as SEQ ID NO:1 of the Sequence Listing.

Please replace the paragraph beginning on page 55, line 18, with the following rewritten paragraph:

The primary structure of the protein (MMR-CAM) (listed as SEQ ID NOS:1 & 2 in the Sequence Listing) deduced from the nucleotide sequence of the gene analyzed (5) consists of 241 amino acid residues, and the molecular weight as estimated from the amino acid sequence was approximately 27 kDa. MMR-CAM is thought to be a type I membrane glycoprotein with one membrane-spanning domain, which comprises an extracellular portion of 107 amino acids, a membrane-spanning portion of 23 amino acids and an intracellular portion of 111 amino acids. Homology search showed that no molecules were found similar to the protein of the invention in terms of the structure, suggesting that the protein of the invention does not belong to the existing family. Also, there is a portion with the extensive modifications by the type O sugar chains was present in the extracellular domain. Phosphorylation sites for protein kinase C, tyrosine kinase, etc. are present in the intracellular domain. These sugar chain binding sites and phosphorylation sites are believed to play on very important role in signal transduction.

Please replace the paragraph beginning on page 56, line 27, with the following rewritten paragraph:

Application No. 09/937,905

A data-search was conducted for human genes homologous to the nucleotide sequence and amino acid sequence listed as SEQ ID NO:1 and determined in Example 1 on both the amino acid level and the DNA level databases using (DNA DATA BANK of JAPAN (DDBJ): Dept. of Education, National Institute of Genetics, Center for Information Biology). The results are shown in Tables 1 and 2. These results suggest that the gene of the invention is also conserved in humans with high homology.

Please replace Table 1 beginning on page 57 with the following rewritten paragraph:

Table 1: Homology on amino acid level

Position within amino acid sequence of SEQ ID NO:1	Matching in human homologue
1 to 91	83/91 (91%)
50 to 146	83/97 (85%)
1 to 78	70/78 (89%)
200 to 241	40/42 (95%)
172 to 241	67/70 (95%)
103 to 150	46/48 (95%)
169 to 241	58/73 (79%)

Please replace Table 2 beginning on page 57 with the following rewritten paragraph:

Table 2: Homology on DNA level

Position within nucleotide sequence of SEQ ID NO:1	Matching in human homologue
519 to 736	189/218 (86%)
666 to 689	23/24 (95%)
381 to 403	22/23 (95%)
709 to 727	19/19 (100%)

Application No. 09/937,905

Please replace the paragraph beginning on page 58, line 3, with the following rewritten paragraph:

Guanidium thiocyanate-phenol-chloroform extraction was used to extract total RNA from human normal brain tissue, and the poly(A)<sup>+</sup> RNA was purified using oligo(dT) cellulose. cDNA was synthesized from the Poly(A)<sup>+</sup> RNA using reverse transcriptase (MMLV-RTase) and DNA polymerase. A sense primer of position 4 to 22 (CCATGTCTGGCTGTCAAGC (SEQ ID NO:5)) and an antisense primer of position 714 to 724 (CCATTTTCTCCAAGTGGGAGC (SEQ ID NO:6)) of the mouse antigen gene (MMR19) sequence were prepared, and these primers and the human normal brain tissue cDNA as the template were used for PCR reaction. As a result, a partial cDNA of the human homologue of the mouse antigen gene (MMR19) was obtained. Next, the 3'RACE method and 5'RACE method were carried out using a specific primer (GSP) for the human homologue partial cDNA and an adapter primer. An antisense primer (GTCAGAAGAGATTTCAGGGTGACC (SEQ ID NO:7)) was prepared from the 3' RACE fragment and a sense primer (AAGCCGTG CGGAGATTGGAGG (SEQ ID NO:8)) from the 5' RACE fragment. As a result of LD-PCR, the full length cDNA of the human homologue including the open reading frame was obtained. The Primer Walking method was used to elucidate the 924 bp nucleotide sequence of the cDNA. The obtained nucleotide sequence is listed as SEQ ID NO:3 of the Sequence Listing. The nucleotide sequence of the human homologue cDNA (924 bp) showed 84.8% homology (with 712 matching nucleotides out of 924) with the nucleotide sequence of the mouse antigen gene cDNA (840 bp).

Please replace the paragraph beginning on page 59, line 1, with the following rewritten paragraph:

The primary structure of the protein deduced from the nucleotide sequence of the obtained gene is listed as SEQ ID NOS:3 & 4, consisting of 242 amino acids. The deduced amino acid sequence showed 93.8% homology with the mouse form (with 226 matching residues out of 242). This protein is also thought to be a type I membrane glycoprotein with one membrane-spanning domain.

Application No. 09/937,905

Please delete the Sequence Listing, originally filed October 1, 2001. Please insert the Substitute Sequence Listing enclosed herewith immediately after the abstract.

*In the Claims:*

1. (Amended) A gene encoding:
  - (a) a protein having the amino acid sequence listed as SEQ ID NO:2 of the Sequence Listing;
  - (b) a protein having the amino acid sequence listed as SEQ ID NO:2 of the Sequence Listing with one or more amino acid deletions, substitutions, additions or insertions and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor; or
  - (c) a protein having at least 50% homology with the amino acid sequence listed as SEQ ID NO:2 and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor.
2. (Amended) A gene having:
  - (a) the nucleotide sequence listed as SEQ ID NO:1 of the Sequence Listing;
  - (b) a nucleotide sequence which is the nucleotide sequence listed as SEQ ID NO:1 of the Sequence Listing with one or more nucleotide deletions, substitutions, additions or insertions and which encodes a protein that can bind to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor; or
  - (c) a nucleotide sequence which hybridizes with DNA having the nucleotide sequence listed as SEQ ID NO:1 of the Sequence Listing under stringent conditions and encodes a protein that can bind to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor.
3. (Amended) A gene encoding:
  - (a) a protein having the amino acid sequence listed as SEQ ID NO:4 of the Sequence Listing;
  - (b) a protein having the amino acid sequence listed as SEQ ID NO:4 of the Sequence Listing with one or more amino acid deletions, substitutions, additions or insertions and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor; or

(c) a protein having at least 50% homology with the amino acid sequence listed as SEQ ID NO:4 and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor.

4. (Amended) A gene having:

(a) the nucleotide sequence listed as SEQ ID NO:3 of the Sequence Listing;

(b) a nucleotide sequence which is the nucleotide sequence listed as SEQ ID NO:3 of the Sequence Listing with one or more nucleotide deletions, substitutions, additions or insertions and which encodes a protein that can bind to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor; or

(c) a nucleotide sequence which hybridizes with DNA having the nucleotide sequence listed as SEQ ID NO:3 of the Sequence Listing under stringent conditions and encodes a protein that can bind to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor.

7. (Amended) A DNA fragment containing:

(1) the nucleotide sequence from position 519 to position 736, the nucleotide sequence from position 666 to position 689, the nucleotide sequence from position 381 to position 403 or the nucleotide sequence from position 709 to position 727 of the nucleotide sequence listed as SEQ ID NO:1 of the Sequence Listing;

(2) a nucleotide sequence which is any of the nucleotide sequences of (1) with one or more nucleotide deletions, substitutions, additions or insertions; or

(3) a nucleotide sequence which has at least 80% homology with any of the nucleotide sequences of (1).

8. (Amended) A gene containing:

(1) the nucleotide sequence from position 519 to position 736, the nucleotide sequence from position 666 to position 689, the nucleotide sequence from position 381 to position 403 or the nucleotide sequence from position 709 to position 727 of the nucleotide sequence listed as SEQ ID NO:1 of the Sequence Listing;

(2) a nucleotide sequence which is any of the nucleotide sequences of (1) with one or more nucleotide deletions, substitutions, additions or insertions; or



(3) a nucleotide sequence which has at least 80% homology with any of the nucleotide sequences of (1),  
and encoding a protein that can bind to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor.

9. (Amended) Any of the following proteins:

- (a) a protein having the amino acid sequence listed as SEQ ID NO:2 of the Sequence Listing;
- (b) a protein having the amino acid sequence listed as SEQ ID NO:2 of the Sequence Listing with one or more amino acid deletions, substitutions, additions or insertions and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor;
- (c) a protein having at least 50% homology with the amino acid sequence listed as SEQ ID NO:2 and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor; or
- (d) a protein that is encoded by the DNA which hybridizes with DNA having the nucleotide sequence listed as SEQ ID NO:1 of the Sequence Listing under stringent conditions and that binds to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor.

10. (Amended) Any of the following proteins:

- (a) a protein having the amino acid sequence listed as SEQ ID NO:4 of the Sequence Listing;
- (b) a protein having the amino acid sequence listed as SEQ ID NO:4 of the Sequence Listing with one or more amino acid deletions, substitutions, additions or insertions and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor;
- (c) a protein having at least 50% homology with the amino acid sequence listed as SEQ ID NO:4 and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor; or
- (d) a protein that is encoded by the DNA which hybridizes with DNA having the nucleotide sequence listed as SEQ ID NO:3 of the Sequence Listing under stringent conditions and that binds to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor.

Application No. 09/937,905

13. (Amended) A protein comprising any of the followings:

(1) the amino acid sequence from residues 1 to 91, the amino acid sequence from residues 50 to 146, the amino acid sequence from residues 1 to 78, the amino acid sequence from residues 200 to 241, the amino acid sequence from residues 172 to 241, the amino acid sequence from residues 103 to 150 or the amino acid sequence from residues 169 to 241 of the amino acid sequence listed as SEQ ID NO:2 of the Sequence Listing;

(2) an amino acid sequence which is any of the amino acid sequences of (1) with one or more amino acid deletions, substitutions, additions or insertions; or

(3) an amino acid sequence having at least 70% homology with any of the amino acid sequences of (1).

14. (Amended) A protein comprising any of the followings:

(1) the amino acid sequence from residues 1 to 91, the amino acid sequence from residues 50 to 146, the amino acid sequence from residues 1 to 78, the amino acid sequence from residues 200 to 241, the amino acid sequence from residues 172 to 241, the amino acid sequence from residues 103 to 150 or the amino acid sequence from residues 169 to 241 of the amino acid sequence listed as SEQ ID NO:2 of the Sequence Listing;

(2) an amino acid sequence which is any of the amino acid sequences of (1) with one or more amino acid deletions, substitutions, additions or insertions; or

(3) an amino acid sequence having at least 70% homology with any of the amino acid sequences of (1), and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor.

*In the Abstract:*

Please replace the Abstract with the rewritten Abstract attached herewith.

Application No. 09/937,905

REMARKS*Sequence Listing*

Enclosed herewith in full compliance with 37 C.F.R. §§1.821-1.825 is a Substitute Sequence Listing to be inserted into the specification as indicated above. The Substitute Sequence Listing in no way introduces new matter into the specification. Also submitted herewith in full compliance with 37 C.F.R. §§1.821-1.825 is a disk copy of the Substitute Sequence Listing. The disk copy of the Substitute Sequence Listing, file "0230-0169P.ST25.TXT", is identical to the paper copy, except that it lacks formatting.

The primer sequences found on page 58 of the Specification as originally filed on October 1, 2001, that were not made part of the Sequence Listing originally filed on same date, are now part of the Substitute Sequence Listing enclosed herewith as SEQ ID NOS: 5 to 8. The Specification has been amended to refer to these primer sequences by their new SEQ ID NOs. In addition, the Specification, Claims, and Abstract have also been amended to replace the references to sequences 1 to 4 from "SEQ. ID. No." to the proper format of "SEQ ID NO". In no way is new matter being added to the application by the addition of the primer sequences to the Sequence Listing nor by the amendments made to the Specification, Claims, and Abstract.

*Claim Fee*

Applicants respectfully request that the requirement to pay the additional claim fee of \$84 as stated in the Notice be withdrawn. The Notice states that the Applicants owe an additional claim fee of \$84 for 8 independent claims over 3. Applicants respectfully point out that there are only 10 (10) independent claims in the present application - claims 1, 2, 3, 4, 7, 8, 9, 10, 13, and 14 - and the required fee of \$588 for filing seven (7) additional independent claims over 3 was paid for at the time of filing. Therefore, since there are only seven (7) additional independent claims over 3, and not eight (8) independent claims over 3, Applicants are not required to pay the additional \$84.

Application No. 09/937,905

If necessary, the Commissioner is hereby authorized in this, concurrent, and future replies, to charge payment or credit any overpayment to Deposit Account No. 02-2448 for any additional fees required under 37 C.F.R. § 1.16 or under 37 C.F.R. § 1.17; particularly, extension of time fees.

Respectfully submitted,  
BIRCH, STEWART, KOLASCH & BIRCH, LLP

By *G. M. Murphy, Jr.*  
Gerald M. Murphy, Jr., Reg. No. 28, 977  
MaryAnne Armstrong, Ph.D., Reg. No. 40,069

*LPS*  
GMM/MAA/LPS

P.O. Box 747  
Falls Church, VA 22040-0747  
(703) 205-8000

## Attachments:

Rewritten Abstract  
Version with Markings Showing Changes Made  
Disk Copy of Substitute Sequence Listing  
Paper Copy of Substitute Sequence Listing  
Copy of Notification

(Rev. 03/27/01)

### **Abstract**

The present invention provides, as a gene encoding an antigen recognized by G-CSF-inducing antibodies, a gene encoding:

(a) a protein having the amino acid sequence listed as SEQ ID NO:2 of the Sequence Listing;

(b) a protein having the amino acid sequence listed as SEQ ID NO:2 of the Sequence Listing with one or more amino acid deletions, substitutions, additions or insertions and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor; or

(c) a protein having at least 50% homology with the amino acid sequence listed as SEQ ID NO:2 and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor.

**VERSION WITH MARKINGS TO SHOW CHANGES MADE**

*In the Specification:*

Please replace the paragraph beginning on page 3, line 17, with the following rewritten paragraph:

In other words, the present invention provides a gene encoding: (a) a protein having the amino acid sequence listed as [SEQ. ID. No.2] **SEQ ID NO:2** of the Sequence Listing; (b) a protein having the amino acid sequence listed as [SEQ. ID. No.2] **SEQ ID NO:2** of the Sequence Listing with one or more amino acid deletions, substitutions, additions or insertions and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor; or (c) a protein having at least 50% homology with the amino acid sequence listed as [SEQ. ID. No.2] **SEQ ID NO:2** and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor.

Please replace the paragraph beginning on page 4, line 1, with the following rewritten paragraph:

The invention further provides a gene encoding: (a) a protein having the amino acid sequence listed as [SEQ. ID. No.4] **SEQ ID NO:4** of the Sequence Listing; (b) a protein having the amino acid sequence listed as [SEQ. ID. No.4] **SEQ ID NO:4** of the Sequence Listing with one or more amino acid deletions, substitutions, additions or insertions and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor; or (c) a protein having at least 50% homology with the amino acid sequence listed as [SEQ. ID. No.4] **SEQ ID NO:4** and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor.

Please replace the paragraph beginning on page 4, line 13, with the following rewritten paragraph:

The invention further provides a gene having: (a) the nucleotide sequence listed as [SEQ. ID. No.1] **SEQ ID NO:1** of the Sequence Listing; (b) a nucleotide sequence which is the nucleotide sequence listed as [SEQ. ID. No.1] **SEQ ID NO:1** of the Sequence Listing with one or more nucleotide deletions, substitutions, additions or insertions and which encodes a protein that can bind to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor; or (c) a nucleotide sequence which hybridizes with DNA having the nucleotide sequence listed as [SEQ. ID. No.1] **SEQ ID NO:1** of the Sequence Listing under stringent conditions and encodes a protein that can bind to an antibody or its fragments that are active to induce granulocyte colony-stimulating factor.

Please replace the paragraph beginning on page 4, line 26, with the following rewritten paragraph:

Application No. 09/937,905

The invention further provides a gene having: (a) the nucleotide sequence listed as [SEQ. ID. No.3] **SEQ ID NO:3** of the Sequence Listing; (b) a nucleotide sequence which is the nucleotide sequence listed as [SEQ. ID. No.3] **SEQ ID NO:3** of the Sequence Listing with one or more nucleotide deletions, substitutions, additions or insertions and which encodes a protein that can bind to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor; or (c) a nucleotide sequence which hybridizes with DNA having the nucleotide sequence listed as [SEQ. ID. No.3] **SEQ ID NO:3** of the Sequence Listing under stringent conditions and encodes a protein that can bind to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor.

Please replace the paragraph beginning on page 5, line 17, with the following rewritten paragraph:

The invention further provides a DNA fragment containing: (1) the nucleotide sequence from position 519 to position 736, the nucleotide sequence from position 666 to position 689, the nucleotide sequence from position 381 to position 403 or the nucleotide sequence from position 709 to position 727 of the nucleotide sequence listed as [SEQ. ID. No.1] **SEQ ID NO:1** of the Sequence Listing; (2) a nucleotide sequence which is any of the nucleotide sequences of (1) above with one or more nucleotide deletions, substitutions, additions or insertions; or (3) a nucleotide sequence which has at least 80% homology with any of the nucleotide sequences of (1) above.

Please replace the paragraph beginning on page 6, line 1, with the following rewritten paragraph:

The invention further provides a gene containing: (1) the nucleotide sequence from position 519 to position 736, the nucleotide sequence from position 666 to position 689, the nucleotide sequence from position 381 to position 403 or the nucleotide sequence from position 709 to position 727 of the nucleotide sequence listed as [SEQ. ID. No.1] **SEQ ID NO:1** of the Sequence Listing; (2) a nucleotide sequence which is any of the nucleotide sequences of (1) above with one or more nucleotide deletions, substitutions, additions or insertions; or (3) a nucleotide sequence which has at least 80% homology with any of the nucleotide sequences of (1) above; and encoding a protein that can bind to an antibody or its fragments that are active to induce granulocyte colony-stimulating factor.

Application No. 09/937,905

Please replace the paragraph beginning on page 6, line 15, with the following rewritten paragraph:

The invention further provides any of the following proteins: (a) a protein having the amino acid sequence listed as [SEQ. ID. No.2] **SEQ ID NO:2** of the Sequence Listing; (b) a protein having the amino acid sequence listed as [SEQ. ID. No.2] **SEQ ID NO:2** of the Sequence Listing with one or more amino acid deletions, substitutions, additions or insertions and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor; (c) a protein having at least 50% homology with the amino acid sequence listed as [SEQ. ID. No.2] **SEQ ID NO:2** and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor; or (d) a protein that is encoded by DNA which hybridizes with DNA having the nucleotide sequence listed as [SEQ. ID. No.1] **SEQ ID NO:1** of the Sequence Listing under stringent conditions and that binds to an antibody or its fragments that are active to induce granulocyte colony-stimulating factor.

Please replace the paragraph beginning on page 7, line 4, with the following rewritten paragraph:

The invention further provides any of the following proteins: (a) a protein having the amino acid sequence listed as [SEQ. ID. No.4] **SEQ ID NO:4** of the Sequence Listing; (b) a protein having the amino acid sequence listed as [SEQ. ID. No.4] **SEQ ID NO:4** of the Sequence Listing with one or more amino acid deletions, substitutions, additions or insertions and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor; (c) a protein having at least 50% homology with the amino acid sequence listed as [SEQ. ID. No.4] **SEQ ID NO:4** and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor; or (d) a protein that is encoded by DNA which hybridizes with DNA having the nucleotide sequence listed as [SEQ. ID. No.3] **SEQ ID NO:3** of the Sequence Listing under stringent conditions and that binds to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor.

Please replace the paragraph beginning on page 7, line 28, with the following rewritten paragraph:

The invention further provides a protein comprising any of the followings: (1) the amino acid sequence from residues 1 to 91, the amino acid sequence from residues 50 to 146, the amino acid sequence from residues 1 to 78, the amino acid sequence from residues 200 to 241, the amino acid



Application No. 09/937,905

sequence from residues 172 to 241, the amino acid sequence from residues 103 to 150 or the amino acid sequence from residues 169 to 241 of the amino acid sequence listed as [SEQ. ID. No.2] **SEQ ID NO:2** of the Sequence Listing; (2) an amino acid sequence which is any of the amino acid sequences of (1) above with one or more amino acid deletions, substitutions, additions or insertions; or (3) an amino acid sequence having at least 70% homology with any of the amino acid sequences of (1) above.

Please replace the paragraph beginning on page 8, line 14, with the following rewritten paragraph:

The invention further provides a protein comprising any of the followings: (1) the amino acid sequence from residues 1 to 91, the amino acid sequence from residues 50 to 146, the amino acid sequence from residues 1 to 78, the amino acid sequence from residues 200 to 241, the amino acid sequence from residues 172 to 241, the amino acid sequence from residues 103 to 150 or the amino acid sequence from residues 169 to 241 of the amino acid sequence listed as [SEQ. ID. No.2] **SEQ ID NO:2** of the Sequence Listing; (2) an amino acid sequence which is any of the amino acid sequences of (1) above with one or more amino acid deletions, substitutions, additions or insertions; or (3) an amino acid sequence having at least 70% homology with any of the amino acid sequences of (1) above, and also binding to an antibody or its fragments that are active to induce granulocyte colony-stimulating factor.

Please replace the paragraph beginning on page 17, line 5, with the following rewritten paragraph:

The present invention provides a gene which encodes the protein having the amino acid sequence listed as [SEQ. ID. No.1] **SEQ ID NO:1** of the Sequence Listing or a protein which is homologous thereto. The invention also provides a gene having the nucleotide sequence listed as [SEQ. ID. No.1] **SEQ ID NO:1** of the Sequence Listing or a nucleotide sequence which is homologous thereto.

Please replace the paragraph beginning on page 17, line 16, with the following rewritten paragraph:

The gene of the invention will typically have the nucleotide sequence listed as [SEQ. ID. No.1] SEQ ID NO:1 of the Sequence Listing, but this is only the sequence of a clone (MMR19) obtained in the following examples to represent an embodiment of the invention. Those skilled in the art are well aware that natural genes have a small number of variations depending on the breeding conditions of the biological species that produce it, and on the ecosystem, or on the presence of highly similar isozymes. Consequently, the "gene" of the invention is not limited to the gene having the nucleotide sequence listed as [SEQ. ID. No.1] SEQ ID NO:1 of the Sequence Listing but also includes all genes encoding proteins having the characters described in the present specification.

Please replace the paragraph beginning on page 19, line 1, with the following rewritten paragraph:

A homologous gene cloned by hybridization in this manner has at least 70%, preferably at least 80%, more preferably at least 90%, even more preferably at least 95% and most preferably at least 98% homology with respect to the nucleotide sequence listed as [SEQ. ID. No.1] SEQ ID NO:1 of the Sequence Listing.

Please replace the paragraph beginning on page 19, line 9, with the following rewritten paragraph:

The present invention provides the protein having the amino acid sequence listed as [SEQ. ID. No.1] SEQ ID NO:1 of the Sequence Listing or a protein homologous thereto.

Please replace the paragraph beginning on page 19, line 12, with the following rewritten paragraph:

The protein having the amino acid sequence listed as [SEQ. ID. No.1] SEQ ID NO:1 of the Sequence Listing according to the invention may be obtained by incorporating the gene encoding therefor into an appropriate expression vector, transforming this vectors to an appropriate host and expressing the recombinant protein. However, the source and preparation method are not restricted so long as the protein of the invention has the characters described in the present specification, and it may be a naturally produced protein, a protein expressed from recombinant DNA by a genetic engineering method or a protein chemically synthesized.

Application No. 09/937,905

Please replace the paragraph beginning on page 19, line 24, with the following rewritten paragraph:

The protein of the invention will typically have the sequence of 241 amino acids listed as [SEQ. ID. No.1] **SEQ ID NO:1** of the Sequence Listing. However, those skilled in the art are well aware that natural proteins include variations of one or more amino acids due to gene variation depending on the breeding [contions] **conditions** of biological species that produce it, on the ecosystem, the presence of highly similar isozymes. The term "amino acid variation" as used here means one or more amino acid substitutions, deletions, insertions and/or additions. The "protein" of the invention has the amino acid sequence listed as [SEQ. ID. No.1] **SEQ ID NO:1** based on deduction from the nucleotide sequence of the cloned gene, but it is not limited only to proteins with that sequence and is intended to include all homologous proteins that have the characteristics described in the present specification. The homology is at least 50%, preferably at least 60%, more preferably at least 70%, more preferably at least 80%, even more preferably at least 90%, yet more preferably at least 95% and most preferably at least 98%.

Please replace the paragraph beginning on page 21, line 16, with the following rewritten paragraph:

The total RNA is extracted from a human macrophage cell line (THP-1, U937, HL-60) by guanidium thiocyanate-phenol-chloroform single-step extraction (Laboratory Manuals of Genetic Engineering, 3rd Edition, pp.83-84, 1996), and purified using an oligo(dT) cellulose column, to obtain poly(A)<sup>+</sup> RNA. Reverse transcriptase (MMLV-RTase) and DNA polymerase are used to synthesize double-stranded cDNA. The double-stranded cDNA is used to construct a cDNA library using a  $\lambda$ ZAPII phage vector by the method of Gubler-Hoffmann (Gubler, U. and Hoffmann, B.J.: Gene, 25:263-269, 1983). A probe is then prepared by amplifying a DNA sequence using a primer DNA that can amplify a sequence in the region of the nucleotide sequence ([SEQ. ID. No.1] **SEQ ID NO:1**) of the mouse cDNA (MMR19 clone) disclosed in the present specification, having high homology with the human sequence (for example, the region from position 172 to position 241 of [SEQ. ID. No.1] **SEQ ID NO:1** which has been found to have 91% homology with the human sequence) and the template DNA from the human macrophage cell cDNA library. Or region (for example, the region from position 172 to position 241 of [SEQ. ID. No.1] **SEQ ID NO:1**) is used

Application No. 09/937,905

directly as the probe for screening the cDNA encoding the entire length of the target protein from the human macrophage cell cDNA library. The cDNA nucleotide sequence is analyzed by the Primer Walking method. The cDNA confirmed to encode the entire length of the target protein is introduced into a baculovirus to express a protein, which can be purified with an affinity column to obtain the human-type homologous protein.

Please replace the paragraph beginning on page 22, line 16, with the following rewritten paragraph:

As explained above, the present invention relates to the gene or protein having the nucleotide sequence listed as [SEQ. ID. No.1] **SEQ ID NO:1** or the amino acid sequence listed as [SEQ. ID. No.2] **SEQ ID NO:2**, and to genes and proteins which are homologous thereto. As a result of a search to determine whether or not sequences homologous to the nucleotide sequence listed as [SEQ. ID. No.1] **SEQ ID NO:1** and the amino acid sequence listed as [SEQ. ID. No.2] **SEQ ID NO:2** provided by the invention are present in other organisms, it was confirmed that human ESTs (expressed sequence tags) include sequences having high homology with the gene of the invention (see Example 3 below). It will therefore be readily appreciated that the human-derived homologous gene can also be isolated by screening of a human-derived gene library (cDNA library, etc.) using the human-derived ESTs having high homology with the nucleotide sequence of the invention as the probe.

Please replace the paragraph beginning on page 23, line 4, with the following rewritten paragraph:

As described above, a database search revealed that portions (i.e. DNA fragments) of the nucleotide sequence listed as [SEQ. ID. No.1] **SEQ ID NO:1** according to the invention are conserved, having high homology with the human sequence. Such DNA fragments are useful as probes for screening of the human-derived homologous gene, and therefore constitute one aspect of the present invention. The DNA fragments include DNA fragments containing the nucleotide sequence from position 519 to position 736, the nucleotide sequence from position 666 to position 689, the nucleotide sequence from position 381 to position 403 or the nucleotide sequence from position 709 to position 727 of the nucleotide sequence listed as [SEQ. ID. No.1] **SEQ ID NO:1** of the Sequence Listing, while DNA fragments containing a nucleotide sequence which is any of these

Application No. 09/937,905

nucleotide sequences with one or more nucleotide deletions, substitutions, additions or insertions or a nucleotide sequence which has at least 80%, preferably at least 85%, more preferably at least 90%, even more preferably at least 95% and most preferably at least 98% homology with any of these nucleotide sequences, are also within the scope of the invention.

Please replace the paragraph beginning on page 23, line 25, with the following rewritten paragraph:

A database search also revealed that a portion of the amino acid sequence listed as [SEQ. ID. No.2] **SEQ ID NO:2** according to the invention is conserved with high homology in the human sequence. Protein fragments comprising portions of the protein of the invention are also useful as reagents for analysis and isolation of antibodies with G-CSF inducing activity, as is the protein of the invention, and also have potential utility as a drug like the protein of the invention, and thus constitute an aspect of the invention.

Please replace the paragraph beginning on page 24, line 6, with the following rewritten paragraph:

The mentioned proteins include the amino acid sequences from residues 1 to 91, 50 to 146, 1 to 78, 200 to 241, 172 to 241, 103 to 150, and 169 to 241 of the amino acid sequence listed as [SEQ. ID. No.2] **SEQ ID NO:2** of the Sequence Listing, while proteins containing an amino acid sequence which is any of these amino acid sequences with one or more amino acid deletions, substitutions, additions or insertions or an amino acid sequence having at least 70%, preferably at least 80%, more preferably at least 85%, even more preferably at least 90%, even more preferably at least 95% and most preferably at least 98% homology with any of these amino acid sequences, are also within the scope of the invention.

Please replace the paragraph beginning on page 24, line 19, with the following rewritten paragraph:

The present inventors have determined the nucleotide sequence of the human-type antigen gene by a method similar to the one described above (see Example 4 below). Consequently, the present invention provides a gene having the nucleotide sequence listed as [SEQ. ID. No.3] **SEQ ID NO:3** of the Sequence Listing or having a nucleotide sequence which is homologous thereto. The

Application No. 09/937,905

invention also provides a protein having the amino acid sequence listed as [SEQ. ID. No.4] **SEQ ID NO:4** of the Sequence Listing or a protein which is homologous thereto. Here, "homologous" means that the scope of the invention is not limited to the gene having the nucleotide sequence listed as [SEQ. ID. No.3] **SEQ ID NO:3** or the protein having the amino acid sequence listed as [SEQ. ID. No.4] **SEQ ID NO:4**, as was explained in the part of "Gene of the invention" and "Protein of the invention" described above.

Please replace the paragraph beginning on page 43, line 19, with the following rewritten paragraph:

The protein of the invention used for screening is preferably (a) a protein having the amino acid sequence listed as [SEQ. ID. No.4] **SEQ ID NO:4** of the Sequence Listing; (b) a protein having the amino acid sequence listed as [SEQ. ID. No.4] **SEQ ID NO:4** of the Sequence Listing with one or more amino acid deletions, substitutions, additions or insertions and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor; (c) a protein having at least 50% (preferably at least 60%, more preferably at least 70%, even more preferably at least 80%, even more preferably at least 90%, especially preferably at least 94% and most preferably at least 98%) homology with the amino acid sequence listed as [SEQ. ID. No.4] **SEQ ID NO:4** and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor; or (d) a protein that is encoded by DNA which hybridizes with DNA having the nucleotide sequence listed as [SEQ. ID. No.3] **SEQ ID NO:3** of the Sequence Listing under stringent conditions and that binds to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor.

Please replace the paragraph beginning on page 55, line 9, with the following rewritten paragraph:

Based on the results of the nucleotide sequence analyses, a clone MMR19 was found to have the 840 bp nucleotide sequence of the full length cDNA which included the open reading frame of the protein. The nucleotide sequence of the clone MMR19 is listed as [SEQ. ID. No.1] **SEQ ID NO:1** of the Sequence Listing.

Please replace the paragraph beginning on page 55, line 18, with the following rewritten paragraph:

Application No. 09/937,905

The primary structure of the protein (MMR-CAM) (listed as [SEQ. ID. No.1 and No.2] **SEQ ID NOS:1 & 2** in the Sequence Listing) deduced from the nucleotide sequence of the gene analyzed (5) consists of 241 amino acid residues, and the molecular weight as estimated from the amino acid sequence was approximately 27 kDa. MMR-CAM is thought to be a type I membrane glycoprotein with one membrane-spanning domain, which comprises an extracellular portion of 107 amino acids, a membrane-spanning portion of 23 amino acids and an intracellular portion of 111 amino acids. Homology search showed that no molecules were found similar to the protein of the invention in terms of the structure, suggesting that the protein of the invention does not belong to the existing family. Also, there is a portion with the extensive modifications by the type O sugar chains was present in the extracellular domain. Phosphorylation sites for protein kinase C, tyrosine kinase, etc. are present in the intracellular domain. These sugar chain binding sites and phosphorylation sites are believed to play on very important role in signal transduction.

Please replace the paragraph beginning on page 56, line 27, with the following rewritten paragraph:

A data-search was conducted for human genes homologous to the nucleotide sequence and amino acid sequence listed as [SEQ. ID. No.1] **SEQ ID NO:1** and determined in Example 1 on both the amino acid level and the DNA level databases using (DNA DATA BANK of JAPAN (DDBJ): Dept. of Education, National Institute of Genetics, Center for Information Biology). The results are shown in Tables 1 and 2. These results suggest that the gene of the invention is also conserved in humans with high homology.

Please replace Table 1 beginning on page 57 with the following rewritten paragraph:

Table 1: Homology on amino acid level

Position within amino acid sequence of [SEQ. ID. No.1] <b>SEQ ID NO:1</b>	Matching in human homologue
1 to 91	83/91 (91%)
50 to 146	83/97 (85%)

Application No. 09/937,905

1 to 78	70/78 (89%)
200 to 241	40/42 (95%)
172 to 241	67/70 (95%)
103 to 150	46/48 (95%)
169 to 241	58/73 (79%)

Please replace Table 2 beginning on page 57 with the following rewritten paragraph:

Table 2: Homology on DNA level

Position within nucleotide sequence of [SEQ. ID. No.1] <b><u>SEQ ID</u></b> <b><u>NO:1</u></b>	Matching in human homologue
519 to 736	189/218 (86%)
666 to 689	23/24 (95%)
381 to 403	22/23 (95%)
709 to 727	19/19 (100%)

Please replace the paragraph beginning on page 58, line 3, with the following rewritten paragraph:

Guanidium thiocyanate-phenol-chloroform extraction was used to extract total RNA from human normal brain tissue, and the poly(A)<sup>+</sup> RNA was purified using oligo(dT) cellulose. cDNA was synthesized from the Poly(A)<sup>+</sup> RNA using reverse transcriptase (MMLV-RTase) and DNA polymerase. A sense primer of position 4 to 22 (CCATGTCTGGCTGTCAAGC (**SEQ ID NO:5**)) and an antisense primer of position 714 to 724 (CCATTTTCTCCAAGTGGGAGC (**SEQ ID NO:6**)) of the mouse antigen gene (MMR19) sequence were prepared, and these primers and the human normal brain tissue cDNA as the template were used for PCR reaction. As a result, a partial cDNA of the human homologue of the mouse antigen gene (MMR19) was obtained. Next, the 3'RACE method and 5'RACE method were carried out using a specific primer (GSP) for the human homologue partial cDNA and an adapter primer. An antisense primer



Application No. 09/937,905

(GTCAGAAGAGATTCAGGGTGACC (**SEQ ID NO:7**)) was prepared from the 3' RACE fragment and a sense primer (AAGCCGTG CGGAGATTGGAGG (**SEQ ID NO:8**)) from the 5' RACE fragment. As a result of LD-PCR, the full length cDNA of the human homologue including the open reading frame was obtained. The Primer Walking method was used to elucidate the 924 bp nucleotide sequence of the cDNA. The obtained nucleotide sequence is listed as [SEQ. ID. No.3] **SEQ ID NO:3** of the Sequence Listing. The nucleotide sequence of the human homologue cDNA (924 bp) showed 84.8% homology (with 712 matching nucleotides out of 924) with the nucleotide sequence of the mouse antigen gene cDNA (840 bp).

Please replace the paragraph beginning on page 59, line 1, with the following rewritten paragraph:

The primary structure of the protein deduced from the nucleotide sequence of the obtained gene is listed as [SEQ. ID. No.3 and No.4] **SEQ ID NOS:3 & 4**, consisting of 242 amino acids. The deduced amino acid sequence showed 93.8% homology with the mouse form (with 226 matching residues out of 242). This protein is also thought to be a type I membrane glycoprotein with one membrane-spanning domain.

*In the Claims:*

1. (Amended) A gene encoding:

(a) a protein having the amino acid sequence listed as [SEQ. ID. No.2] **SEQ ID NO:2** of the Sequence Listing;

(b) a protein having the amino acid sequence listed as [SEQ. ID. No.2] **SEQ ID NO:2** of the Sequence Listing with one or more amino acid deletions, substitutions, additions or insertions and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor; or

(c) a protein having at least 50% homology with the amino acid sequence listed as [SEQ. ID. No.2] **SEQ ID NO:2** and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor.

2. (Amended) A gene having:
  - (a) the nucleotide sequence listed as [SEQ. ID. No.1] **SEQ ID NO:1** of the Sequence Listing;
  - (b) a nucleotide sequence which is the nucleotide sequence listed as [SEQ. ID. No.1] **SEQ ID NO:1** of the Sequence Listing with one or more nucleotide deletions, substitutions, additions or insertions and which encodes a protein that can bind to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor; or
  - (c) a nucleotide sequence which hybridizes with DNA having the nucleotide sequence listed as [SEQ. ID. No.1] **SEQ ID NO:1** of the Sequence Listing under stringent conditions and encodes a protein that can bind to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor.
3. (Amended) A gene encoding:
  - (a) a protein having the amino acid sequence listed as [SEQ. ID. No.4] **SEQ ID NO:4** of the Sequence Listing;
  - (b) a protein having the amino acid sequence listed as [SEQ. ID. No.4] **SEQ ID NO:4** of the Sequence Listing with one or more amino acid deletions, substitutions, additions or insertions and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor; or
  - (c) a protein having at least 50% homology with the amino acid sequence listed as [SEQ. ID. No.4] **SEQ ID NO:4** and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor.
4. (Amended) A gene having:
  - (a) the nucleotide sequence listed as [SEQ. ID. No.3] **SEQ ID NO:3** of the Sequence Listing;
  - (b) a nucleotide sequence which is the nucleotide sequence listed as [SEQ. ID. No.3] **SEQ ID NO:3** of the Sequence Listing with one or more nucleotide deletions, substitutions, additions or insertions and which encodes a protein that can bind to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor; or
  - (c) a nucleotide sequence which hybridizes with DNA having the nucleotide sequence listed as [SEQ. ID. No.3] **SEQ ID NO:3** of the Sequence Listing under stringent conditions and encodes a

protein that can bind to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor.

7. (Amended) A DNA fragment containing:

(1) the nucleotide sequence from position 519 to position 736, the nucleotide sequence from position 666 to position 689, the nucleotide sequence from position 381 to position 403 or the nucleotide sequence from position 709 to position 727 of the nucleotide sequence listed as [SEQ. ID. No.1] **SEQ ID NO:1** of the Sequence Listing;

(2) a nucleotide sequence which is any of the nucleotide sequences of (1) with one or more nucleotide deletions, substitutions, additions or insertions; or

(3) a nucleotide sequence which has at least 80% homology with any of the nucleotide sequences of (1).

8. (Amended) A gene containing:

(1) the nucleotide sequence from position 519 to position 736, the nucleotide sequence from position 666 to position 689, the nucleotide sequence from position 381 to position 403 or the nucleotide sequence from position 709 to position 727 of the nucleotide sequence listed as [SEQ. ID. No.1] **SEQ ID NO:1** of the Sequence Listing;

(2) a nucleotide sequence which is any of the nucleotide sequences of (1) with one or more nucleotide deletions, substitutions, additions or insertions; or

(3) a nucleotide sequence which has at least 80% homology with any of the nucleotide sequences of (1),

and encoding a protein that can bind to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor.

9. (Amended) Any of the following proteins:

(a) a protein having the amino acid sequence listed as [SEQ. ID. No.2] **SEQ ID NO:2** of the Sequence Listing;

(b) a protein having the amino acid sequence listed as [SEQ. ID. No.2] **SEQ ID NO:2** of the Sequence Listing with one or more amino acid deletions, substitutions, additions or insertions and

Application No. 09/937,905

also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor;

(c) a protein having at least 50% homology with the amino acid sequence listed as [SEQ. ID. No.2] **SEQ ID NO:2** and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor; or

(d) a protein that is encoded by the DNA which hybridizes with DNA having the nucleotide sequence listed as [SEQ. ID. No.1] **SEQ ID NO:1** of the Sequence Listing under stringent conditions and that binds to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor.

10. (Amended) Any of the following proteins:

(a) a protein having the amino acid sequence listed as [SEQ. ID. No.4] **SEQ ID NO:4** of the Sequence Listing;

(b) a protein having the amino acid sequence listed as [SEQ. ID. No.4] **SEQ ID NO:4** of the Sequence Listing with one or more amino acid deletions, substitutions, additions or insertions and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor;

(c) a protein having at least 50% homology with the amino acid sequence listed as [SEQ. ID. No.4] **SEQ ID NO:4** and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor; or

(d) a protein that is encoded by the DNA which hybridizes with DNA having the nucleotide sequence listed as [SEQ. ID. No.3] **SEQ ID NO:3** of the Sequence Listing under stringent conditions and that binds to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor.

[ 113. (Amended) A protein comprising any of the followings:

(1) the amino acid sequence from residues 1 to 91, the amino acid sequence from residues 50 to 146, the amino acid sequence from residues 1 to 78, the amino acid sequence from residues 200 to 241, the amino acid sequence from residues 172 to 241, the amino acid sequence from residues 103 to 150 or the amino acid sequence from residues 169 to 241 of the amino acid sequence listed as [SEQ. ID. No.2] **SEQ ID NO:2** of the Sequence Listing;

Application No. 09/937,905

(2) an amino acid sequence which is any of the amino acid sequences of (1) with one or more amino acid deletions, substitutions, additions or insertions; or

(3) an amino acid sequence having at least 70% homology with any of the amino acid sequences of (1).

14. (Amended) A protein comprising any of the followings:

(1) the amino acid sequence from residues 1 to 91, the amino acid sequence from residues 50 to 146, the amino acid sequence from residues 1 to 78, the amino acid sequence from residues 200 to 241, the amino acid sequence from residues 172 to 241, the amino acid sequence from residues 103 to 150 or the amino acid sequence from residues 169 to 241 of the amino acid sequence listed as [SEQ. ID. No.2] **SEQ ID NO:2** of the Sequence Listing;

(2) an amino acid sequence which is any of the amino acid sequences of (1) with one or more amino acid deletions, substitutions, additions or insertions; or

(3) an amino acid sequence having at least 70% homology with any of the amino acid sequences of (1), and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor.

Application No. 09/937,905

Abstract

The present invention provides, as a gene encoding an antigen recognized by G-CSF-inducing antibodies, a gene encoding:

(a) a protein having the amino acid sequence listed as [SEQ. ID. No.2]**SEQ ID NO:2** of the Sequence Listing;

(b) a protein having the amino acid sequence listed as [SEQ. ID. No.2]**SEQ ID NO:2** of the Sequence Listing with one or more amino acid deletions, substitutions, additions or insertions and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor; or

(c) a protein having at least 50% homology with the amino acid sequence listed as [SEQ. ID. No.2]**SEQ ID NO:2** and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor.

## SEQUENCE LISTING

<110> SHA, Shiken  
 AOKI, Yoshiko  
 NISHI, Yoshisuke

<120> Novel Proteins, Gene Encoding the Same and Method of Utilization Thereof

<130> 0230-0169P

<140> US 09/937,905  
 <141> 2001-10-01

<150> PCT/JP00/02080  
 <151> 2000-03-31

<150> Japan 95092/1999  
 <151> 1999-04-01

<160> 8

<170> PatentIn version 3.1

<210> 1  
 <211> 840  
 <212> DNA  
 <213> Mus sp.

<220>  
 <221> CDS  
 <222> (6)..(731)  
 <223>

<220>  
 <221> misc\_feature  
 <222> (1)..(840)  
 <223> Mouse macrophage cell RAW 264.7

<400> 1  
 gaacc atg tct ggc tgt caa gct caa gga gac tgt tgc tgc cgg ccg tgt 50  
 Met Ser Gly Cys Gln Ala Gln Gly Asp Cys Cys Ser Arg Pro Cys  
 1 5 10 15  
 ggc gcg cag gac aag gag cac ccc cga ttc ctg atc cca gaa ctt tgc 98  
 Gly Ala Gln Asp Lys Glu His Pro Arg Phe Leu Ile Pro Glu Leu Cys  
 20 25 30  
 aaa cag ttt tac cat ctg ggc tgg gtc act ggc act gga ggg gga atc 146  
 Lys Gln Phe Tyr His Leu Gly Trp Val Thr Gly Thr Gly Gly Gly Ile  
 35 40 45  
 agc ttg aag cat ggc aat gaa atc tac att gct ccc tca ggc gtg caa 194  
 Ser Leu Lys His Gly Asn Glu Ile Tyr Ile Ala Pro Ser Gly Val Gln  
 50 55 60







195

200

205

Met Cys Glu Cys Tyr Asp Tyr Leu Phe Asp Ile Ala Val Ser Met Lys  
210 215 220

Lys Met Gly Leu Asp Pro Thr Gln Leu Pro Val Gly Glu Asn Gly Ile  
225 230 235 240

Val

<210> 3  
<211> 924  
<212> DNA  
<213> Homo sapiens

<220>  
<221> CDS  
<222> (46)..(774)  
<223>

<220>  
<221> misc\_feature  
<222> (1)..(924)  
<223> Human normal brain tissue

<400> 3  
aagccgtgcg gagattggag gccgcgcggg tccctggtct gggcc atg tct ggc tgt 57  
Met Ser Gly Cys  
1  
  
gat gct tgg gag gga gac tgt tgt tcc cgg aga tgc ggc gcg cag gac 105  
Asp Ala Trp Glu Gly Asp Cys Cys Ser Arg Arg Cys Gly Ala Gln Asp  
5 10 15 20  
  
aag gag cat cca aga tac ctg atc cca gaa ctt tgc aaa cag ttt tac 153  
Lys Glu His Pro Arg Tyr Leu Ile Pro Glu Leu Cys Lys Gln Phe Tyr  
25 30 35  
  
cat tta ggc tgg gtc act ggg act gga gga gga att agc ttg aag cat 201  
His Leu Gly Trp Val Thr Gly Thr Gly Gly Gly Ile Ser Leu Lys His  
40 45 50  
  
ggc gat gaa atc tac att gct cct tca gga gtg caa aag gaa cga att 249  
Gly Asp Glu Ile Tyr Ile Ala Pro Ser Gly Val Gln Lys Glu Arg Ile  
55 60 65  
  
cag cct gaa gac atg ttt gtt tat gat ata aat gaa aag gac ata agt 297  
Gln Pro Glu Asp Met Phe Val Tyr Asp Ile Asn Glu Lys Asp Ile Ser  
70 75 80

```

gga cct tcg cca tcg aag aag cta aaa aaa agc cag tgt act cct ctt      345
Gly Pro Ser Pro Ser Lys Lys Leu Lys Lys Ser Gln Cys Thr Pro Leu
85                      90                      95                      100

ttc atg aat gct tac aca atg aga gga gca ggt gca gtg att cat acc      393
Phe Met Asn Ala Tyr Thr Met Arg Gly Ala Gly Ala Val Ile His Thr
                      105                      110                      115

cac tct aaa gct gct gtg atg gcc acc ctt ctc ttt cca gga cgg gag      441
His Ser Lys Ala Ala Val Met Ala Thr Leu Leu Phe Pro Gly Arg Glu
                      120                      125                      130

ttt aaa att aca cat caa gag atg ata aaa gga ata aag aaa tgt act      489
Phe Lys Ile Thr His Gln Glu Met Ile Lys Gly Ile Lys Lys Cys Thr
                      135                      140                      145

tcc gga ggg tat tat aga tat gat gat atg tta gtg gta ccc att att      537
Ser Gly Gly Tyr Tyr Arg Tyr Asp Asp Met Leu Val Val Pro Ile Ile
                      150                      155                      160

gag aat aca cct gag gag aaa gac ctc aaa gat aga atg gct cat gca      585
Glu Asn Thr Pro Glu Glu Lys Asp Leu Lys Asp Arg Met Ala His Ala
165                      170                      175                      180

atg aat gaa tac cca gac tcc tgt gca gta ctg gtc aga cgt cat gga      633
Met Asn Glu Tyr Pro Asp Ser Cys Ala Val Leu Val Arg Arg His Gly
                      185                      190                      195

gta tat gtg tgg ggg gaa aca tgg gag aag gcc aaa acc atg tgt gag      681
Val Tyr Val Trp Gly Glu Thr Trp Glu Lys Ala Lys Thr Met Cys Glu
                      200                      205                      210

tgt tat gac tat tta ttt gat att gcc gta tca atg aag aaa gta gga      729
Cys Tyr Asp Tyr Leu Phe Asp Ile Ala Val Ser Met Lys Lys Val Gly
                      215                      220                      225

ctt gat cct tca cag ctc cca gtt gga gaa aat gga att gtc taa      774
Leu Asp Pro Ser Gln Leu Pro Val Gly Glu Asn Gly Ile Val
                      230                      235                      240

gccaaaagaa agtctaatta tatacagaga taaagctaaa cgtaattatt atttaaata      834
aagctatttt tttaaatagaa ttgaaatttt tcatgatgct actaatttgc cactaaatac      894
tgcaaatggg caccctgaat ctcttctgac      924

```

```

<210> 4
<211> 242
<212> PRT
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<222> (1)..(242)
<223> Human normal brain tissue

```

&lt;400&gt; 4

Met Ser Gly Cys Asp Ala Trp Glu Gly Asp Cys Cys Ser Arg Arg Cys  
1 5 10 15

Gly Ala Gln Asp Lys Glu His Pro Arg Tyr Leu Ile Pro Glu Leu Cys  
20 25 30

Lys Gln Phe Tyr His Leu Gly Trp Val Thr Gly Thr Gly Gly Gly Ile  
35 40 45

Ser Leu Lys His Gly Asp Glu Ile Tyr Ile Ala Pro Ser Gly Val Gln  
50 55 60

Lys Glu Arg Ile Gln Pro Glu Asp Met Phe Val Tyr Asp Ile Asn Glu  
65 70 75 80

Lys Asp Ile Ser Gly Pro Ser Pro Ser Lys Lys Leu Lys Lys Ser Gln  
85 90 95

Cys Thr Pro Leu Phe Met Asn Ala Tyr Thr Met Arg Gly Ala Gly Ala  
100 105 110

Val Ile His Thr His Ser Lys Ala Ala Val Met Ala Thr Leu Leu Phe  
115 120 125

Pro Gly Arg Glu Phe Lys Ile Thr His Gln Glu Met Ile Lys Gly Ile  
130 135 140

Lys Lys Cys Thr Ser Gly Gly Tyr Tyr Arg Tyr Asp Asp Met Leu Val  
145 150 155 160

Val Pro Ile Ile Glu Asn Thr Pro Glu Glu Lys Asp Leu Lys Asp Arg  
165 170 175

Met Ala His Ala Met Asn Glu Tyr Pro Asp Ser Cys Ala Val Leu Val  
180 185 190

Arg Arg His Gly Val Tyr Val Trp Gly Glu Thr Trp Glu Lys Ala Lys  
195 200 205

Thr Met Cys Glu Cys Tyr Asp Tyr Leu Phe Asp Ile Ala Val Ser Met  
210 215 220

Lys Lys Val Gly Leu Asp Pro Ser Gln Leu Pro Val Gly Glu Asn Gly  
 225                                    230                                    235                                    240

Ile Val

<210> 5  
 <211> 19  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Sense primer of position 4 to 22 of the mouse antigen gene MMR19

<400> 5  
 ccatgtctgg ctgtcaagc 19

<210> 6  
 <211> 21  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> An antisense primer of position 714 to 724 of the mouse anitgen g  
 ene MMR19

<400> 6  
 ccattttctc caactgggag c 21

<210> 7  
 <211> 23  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Antisense primer of the 3' RACE fragment

<400> 7  
 gtcagaagag attcaggggtg acc 23

<210> 8  
 <211> 21  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Sense primer of the 5' RACE fragment

<400> 8  
 aagccgtgcg gagattggag g 21



UNITED STATES PATENT AND TRADEMARK OFFICE

Commissioner for Patents, Box PCT  
United States Patent and Trademark Office  
Washington, D.C. 20231  
www.uspto.gov

U.S. APPLICATION NUMBER NO.	FIRST NAMED APPLICANT	ATTY. DOCKET NO.
09/937,905	Shiken Sha	0230-0169P

INTERNATIONAL APPLICATION NO.
-------------------------------

PCT/JP00/02080


I.A. FILING DATE	PRIORITY DATE
03/31/2000	04/01/1999

2292  
BIRCH STEWART KOLASCH & BIRCH  
PO BOX 747  
FALLS CHURCH, VA 22040-0747

*mm*  
**Docketed**  
**1-14-02**

CONFIRMATION NO. 5513

371 FORMALITIES LETTER



\*OC000000007081909\*

Date Mailed: 11/14/2001

*Sequence + claim*  
*Listing fee*

### NOTIFICATION OF MISSING REQUIREMENTS UNDER 35 U.S.C. 371 IN THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US)

The following items have been submitted by the applicant or the IB to the United States Patent and Trademark Office as an Elected Office (37 CFR 1.495):

- U.S. Basic National Fees
- Priority Document
- Biochemical Sequence Diskette
- Biochemical Sequence Listing
- Copy of IPE Report
- Copy of references cited in ISR
- Copy of the International Application
- Copy of the International Search Report
- Information Disclosure Statements
- Oath or Declaration
- Preliminary Amendments
- Request for Immediate Examination

The following items **MUST** be furnished within the period set forth below in order to complete the requirements for acceptance under 35 U.S.C. 371:

- The nucleotide and/or amino acid sequence disclosure contained in this application does not comply with the requirements for such a disclosure as set forth in 37 CFR 1.821-1.825 for the following reason(s):
  - NO DISKETTE
  - APPLICANT MUST PROVIDE:
    - An initial or substitute paper copy or compact disc of the "Sequence Listing," as well as an amendment directing its entry into the specification.

- A statement that the contents of the paper or compact disc and the computer readable form are the same and, where applicable, include no new matter, as required by 37 CFR 1.821(e), 1.821(f), 1.821(g), 1.825(b) or 1.825(d).
- For questions regarding compliance to 37 CFR 1.821-1.825 requirements, please contact:
  - For Rules Interpretation, call (703) 308-4216
  - To Purchase PatentIn Software, call (703) 306-2600
  - For PatentIn Software Program Help, call (703) 306-4119 or e-mail at [patin21help@uspto.gov](mailto:patin21help@uspto.gov) or [patin3help@uspto.gov](mailto:patin3help@uspto.gov)

ALL OF THE ITEMS SET FORTH ABOVE MUST BE SUBMITTED WITHIN TWO (2) MONTH FROM THE DATE OF THIS NOTICE OR BY 22 or 32 MONTHS (where 37 CFR 1.495 applies) FROM THE PRIORITY DATE FOR THE APPLICATION, WHICHEVER IS LATER. FAILURE TO PROPERLY RESPOND WILL RESULT IN ABANDONMENT.

The time period set above may be extended by filing a petition and fee for extension of time under the provisions of 37 CFR 1.136(a).

Additionally the following defects have been observed:

- Additional claim fees of **\$84** as a non-small entity, including any required multiple dependent claim fee, are required. Applicant must submit the additional claim fees or cancel the additional claims for which fees are due.

SUMMARY OF FEES DUE:

Total additional fees required for this application is **\$84** for a Large Entity:

- Total additional claim fee(s) for this application is **\$84**
  - **\$84** for **8** independent claims over 3.

Applicant is reminded that any communications to the United States Patent and Trademark Office must be mailed to the address given in the heading and include the U.S. application no. shown above (37 CFR 1.5)

*A copy of this notice **MUST** be returned with the response.*

TAMALA D HOLLAND

Telephone: (703) 305-5483

PART 1 - ATTORNEY/APPLICANT COPY

U.S. APPLICATION NUMBER NO.	INTERNATIONAL APPLICATION NO.	ATTY. DOCKET NO.
09/937,905	PCT/JP00/02080	0230-0169P

09/937905

09/937905

JC05 Rec'd PCT/PTO 0 1 OCT 2001

PATENT  
0230-0169P

IN THE U.S. PATENT AND TRADEMARK OFFICE

Applicant: SHA, Shiken et al. Conf.:  
Int'l. Appl. No.: PCT/JP00/02080  
Appl. No.: New Group:  
Filed: October 1, 2001 Examiner:  
For: NOVEL PROTEINS, GENE ENCODING THE SAME AND  
METHOD OF UTILIZATION THEREOF

PRELIMINARY AMENDMENT

**BOX PATENT APPLICATION**

Assistant Commissioner for Patents  
Washington, DC 20231

October 1, 2001

Sir:

The following Preliminary Amendments and Remarks are respectfully submitted in connection with the above-identified application.

AMENDMENTS

IN THE SPECIFICATION:

Please amend the specification as follows:

Before line 1, insert --This application is the national phase under 35 U.S.C. § 371 of PCT International Application No. PCT/JP00/02080 which has an International filing date of March 31, 2000, which designated the United States of America.--



Docket No. 0230-0169P

IN THE CLAIMS:

Please amend the claims as follows:

6. (Amended) A gene according to claim 1, which is a mouse-derived or human derived gene.

12. (Amended) A protein according to claim 9, which is a mouse-derived, human-derived or other mammalian-derived protein.

15. (Amended) An antibody against a protein according to claim 9, or a fragment thereof.

18. (Amended) A recombinant vector containing a gene or DNA fragment according to claim 1.

19. (Amended) A transformant comprising a recombinant vector that contains the gene or the DNA fragment according to claim 1.

20. (Amended) A receptor for a substance that can induce production of granulocyte colony-stimulating factor including, such as a monoclonal antibody or its fragment that is produced by a hydridoma of the cell line deposited as FERM BP-6103, and the receptor comprises a protein according to claim 9 and is present in a cell which can produce granulocyte colony-stimulating factor, such as macrophage.

21. (Amended) A screening method for any of the following

substances, the method includes measurement of binding between a substance and a protein according to claim 9 or a receptor according to claim 20, measurement of the effect of the substance via a receptor according to claim 20, or measurement to compare the effect between the structure of a substance and the structure of a protein according to the invention.

- (a) a substance which can bind to a receptor according to claim 20, and as a result of its binding to a receptor, it can cause a change in the receptor structure, transmit signals into the cell via the receptor, and induce production of granulocyte colony-stimulating factor;
- (b) a substance which can bind to a receptor according to claim 20, and as a result of its binding to a receptor, it can inhibit the binding of the receptor to the substances that can induce production of granulocyte colony-stimulating factor, but it in itself does not induce production of granulocyte colony-stimulating factor; or
- (c) a substance which can bind to a receptor according to claim 20, and as a result of its binding to a receptor, it can inhibit the binding of the receptor to the substances that can induce production of granulocyte colony-stimulating factor, but it in itself blocks production of granulocyte colony-stimulating factor.

22. (Amended) Any of the following substances obtained by a

screening method, which is characterized by measurement of binding between a substance and a protein according to claim 9 or a receptor according to claim 20, or measurement of the effect of the substance via a receptor according to claim 20,

- (a) a substance which can bind to a receptor according to claim 20, and as a result of its binding to a receptor, it can cause a change in the receptor structure, transmit signals into the cell via the receptor, and induce production of granulocyte colony-stimulating factor;
- (b) a substance which can bind to a receptor according to claim 20, and as a result of its binding to a receptor, it can inhibit the binding of the receptor to the substances that can induce production of granulocyte colony-stimulating factor, but it in itself does not induce production of granulocyte colony-stimulating factor; or
- (c) a substance which can bind to a receptor according to claim 20, and as result of its binding to a receptor, it can inhibit the binding of the receptor to the substances that can induce production of granulocyte colony-stimulating factor, but it in itself blocks production of granulocyte colony-stimulating factor.

24. (Amended) A pharmaceutical composition comprising a gene or DNA fragment according to claim 1, a protein according to claim 9, an antibody or its fragment according to claim 15, a



REMARKS

The specification has been amended to provide a cross-reference to the previously filed International Application. The claims have also been amended to delete improper multiple dependencies and to place the application into better form for examination. Entry of the present amendment and favorable action on the above-identified application are earnestly solicited.

Attached hereto is a marked-up copy of the changes made to the application by this Amendment.

If necessary, the Commissioner is hereby authorized in this, concurrent, and future replies, to charge payment or credit any overpayment to Deposit Account No. 02-2448 for any additional fees required under 37 C.F.R. § 1.16 or under 37 C.F.R. § 1.17; particularly, extension of time fees.

Respectfully submitted,

BIRCH, STEWART, KOLASCH & BIRCH, LLP

By *Gerald M. Murphy, Jr.* #36,623  
Gerald M. Murphy, Jr., #28,977

GMM/cqc  
0230-0169P

P.O. Box 747  
Falls Church, VA 22040-0747  
(703) 205-8000

Attachment: Version With Markings Showing Changes Made

Docket No. 0230-0169P

**VERSION WITH MARKINGS SHOWING CHANGES MADE**

The specification has been amended to provide cross-referencing to the International Application.

The claims have been amended as follows:

6. (Amended) A gene according to [any one of claims 1 to 5]claim 1, which is a mouse-derived or human derived gene.

12. (Amended) A protein according to [any one of claims 9 to 11]claim 9, which is a mouse-derived, human-derived or other mammalian-derived protein.

15. (Amended) An antibody against a protein according to [any one of claims 9 to 14]claim 9, or a fragment thereof.

18. (Amended) A recombinant vector containing a gene or DNA fragment according to [any one of claims 1 to 8]claim 1.

19. (Amended) A transformant comprising a recombinant vector that contains the gene or the DNA fragment according to [any one of claims 1 to 8]claim 1.

20. (Amended) A receptor for a substance that can induce production of granulocyte colony-stimulating factor including, such as a monoclonal antibody or its fragment that is produced by a hydridoma of the cell line deposited as FERM BP-6103, and the

Docket No. 0230-0169P

receptor comprises a protein according to [any one of claims 9 to 12]claim 9 and is present in a cell which can produce granulocyte colony-stimulating factor, such as macrophage.

21. (Amended) A screening method for any of the following substances, the method includes measurement of binding between a substance and a protein according to [any one of claims 9 to 12]claim 9 or a receptor according to claim 20, measurement of the effect of the substance via a receptor according to claim 20, or measurement to compare the effect between the structure of a substance and the structure of a protein according to the invention,

- (a) a substance which can bind to a receptor according to claim 20, and as a result of its binding to a receptor, it can cause a change in the receptor structure, transmit signals into the cell via the receptor, and induce production of granulocyte colony-stimulating factor;
- (b) a substance which can bind to a receptor according to claim 20, and as a result of its binding to a receptor, it can inhibit the binding of the receptor to the substances that can induce production of granulocyte colony-stimulating factor, but it in itself does not induce production of granulocyte colony-stimulating factor; or
- (c) a substance which can bind to a receptor according to claim 20, and as a result of its binding to a receptor,

Docket No. 0230-0169P

it can inhibit the binding of the receptor to the substances that can induce production of granulocyte colony-stimulating factor, but it in itself blocks production of granulocyte colony-stimulating factor.

22. (Amended) Any of the following substances obtained by a screening method, which is characterized by measurement of binding between a substance and a protein according to [any one of claims 9 to 12]claim 9 or a receptor according to claim 20, or measurement of the effect of the substance via a receptor according to claim 20,

- (a) a substance which can bind to a receptor according to claim 20, and as a result of its binding to a receptor, it can cause a change in the receptor structure, transmit signals into the cell via the receptor, and induce production of granulocyte colony-stimulating factor;
- (b) a substance which can bind to a receptor according to claim 20, and as a result of its binding to a receptor, it can inhibit the binding of the receptor to the substances that can induce production of granulocyte colony-stimulating factor, but it in itself does not induce production of granulocyte colony-stimulating factor; or
- (c) a substance which can bind to a receptor according to claim 20, and as result of its binding to a receptor, it can inhibit the binding of the receptor to the



Docket No. 0230-0169P

substances that can induce production of granulocyte colony-stimulating factor, but it in itself blocks production of granulocyte colony-stimulating factor.

24. (Amended) A pharmaceutical composition comprising a gene or DNA fragment according to [any one of claims 1 to 8]claim 1, a protein according to [any one of claims 9 to 14]claim 9, an antibody or its fragment according to [any one of claims 15 to 17]claim 15, a receptor according to claim 20 or a substance according to claim [22 or]23.

26. (Amended) A method of diagnosis, prevention or treatment of a disease or a condition related to G-CSF, such as infection or neutropenia, which employs a gene or DNA fragment according to [any one of claims 1 to 8]claim 1, a protein according to [any one of claims 9 to 14]claim 9, an antibody or its fragment according to [any one of claims 15 to 17]claim 15, a receptor according to claim 20 or a substance according to claim [22 or]23.

27. (Amended) The use of a gene or DNA fragment according to [any one of claims 1 to 8]claim 1, a protein according to [any one of claims 9 to 14]claim 9, an antibody or its fragment according to [any one of claims 15 to 17]claim 15, a receptor according to claim 20 or a substance according to claim [22 or]23, for diagnosis, prevention or treatment of a disease or condition related to G-CSF, such as infection or neutropenia.

Docket No. 0230-0169P

28. (Amended) The use of a gene or DNA fragment according to [any one of claims 1 to 8]claim 1, a protein according to [any one of claims 9 to 14]claim 9, an antibody or its fragment according to [any one of claims 15 to 17]claim 15, a receptor according to claim 20 or a substance according to claim [22 or]23, in the production of a drug for diagnosis, prevention or treatment of a disease or condition related to G-CSF, such as infection or neutropenia.

## SPECIFICATION

NOVEL PROTEINS, GENE ENCODING THE SAME  
AND METHOD OF UTILIZATION THEREOF

5

TECHNICAL FIELD

The present invention relates to a protein which is reactive with antibodies that are active to induce granulocyte colony-stimulating factor, to a gene encoding  
10 it and to a method for their use.

BACKGROUND ART

Granulocyte colony-stimulating factor (G-CSF) has a molecular weight of approximately 18,000 to 22,000 and  
15 consists of 174 (in rare cases 178) amino acids in the case of humans and 178 amino acids in the case of mice. It is a glycoprotein that induces differentiation and proliferation of neutrophils, one of the types of leukocytes.

G-CSF has a potential of survival-extension and  
20 functional promotion to the mature neutrophils, and also has ability to form erythroblasts in response to erythropoietin and blast cell colonies in response to interleukin-3. Cells that produce G-CSF are macrophages, stroma cells, monocytes, T lymphocytes, fibroblasts,  
25 vascular endothelial cells and so forth.

Administration of G-CSF drug exhibits a therapeutic effect on neutropenia induced by side effect of anticancer agents, or neutropenia following bone marrow

transplantation, and a therapeutic effect on anaplastic anemia. Because of its low stability in the blood, however, it requires frequent administration, and because its administration is limited to the intravenous route, this  
5 has resulted in a great deal of pain and burden to the patient and physician. Furthermore, administration of G-CSF as a drug has been reported to cause ostealgia as a side-effect. The alternative option of direct administration of macrophages or stroma cells that produce G-CSF  
10 will produce the risk of unknown side-effects since the cells contain numerous proteins and other substances, and therefore such treatment has not been practiced.

Because administration of G-CSF itself for differentiation and proliferation of neutrophils provoke  
15 ostealgia as a side-effect, and it also requires frequent administration and increases the pain and burden to the patient and physician, it has been strongly desired to develop an alternative treatment method; however, no such method has yet been established.

20 With the intent of causing production of G-CSF and differentiation and proliferation of neutrophils without administration of G-CSF itself, the present inventors have already succeeded in providing G-CSF inducing antibodies (Japanese Patent Application HEI No. 9-266591 (Sept. 30,  
25 1997), Japanese Unexamined Patent Publication HEI No. 11-106400 (April 20, 1999)).

However, the antigens recognized by the G-CSF inducing antibodies have not yet been discovered.

One problem to be solved by the present invention, therefore, is to identify an antigen recognized by G-CSF inducing antibodies. Another problem to be solved by the invention is to clone and identify the gene encoding the antigen recognized by the G-CSF inducing antibodies.

#### DISCLOSURE OF THE INVENTION

As a result of diligent research aimed at solving the problems described above, the present inventors used monoclonal antibodies with G-CSF inducing ability as probes for immunoscreening of a cDNA library derived from macrophage cells, and as a result succeeded in isolating 3 positive clones, and then further determined the nucleotide sequences thereof, to thus arrive at the present invention. The present inventors also determined the nucleotide sequence of the human antigen gene.

In other words, the present invention provides a gene encoding: (a) a protein having the amino acid sequence listed as SEQ. ID. No.2 of the Sequence Listing; (b) a protein having the amino acid sequence listed as SEQ. ID. No.2 of the Sequence Listing with one or more amino acid deletions, substitutions, additions or insertions and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor; or (c) a protein having at least 50% homology with the amino acid sequence listed as SEQ. ID. No.2 and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor.

The invention further provides a gene encoding: (a) a protein having the amino acid sequence listed as SEQ. ID. No.4 of the Sequence Listing; (b) a protein having the amino acid sequence listed as SEQ. ID. No.4 of the Sequence Listing with one or more amino acid deletions, substitutions, additions or insertions and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor; or (c) a protein having at least 50% homology with the amino acid sequence listed as SEQ. ID. No.4 and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor.

The invention further provides a gene having: (a) the nucleotide sequence listed as SEQ. ID. No.1 of the Sequence Listing; (b) a nucleotide sequence which is the nucleotide sequence listed as SEQ. ID. No.1 of the Sequence Listing with one or more nucleotide deletions, substitutions, additions or insertions and which encodes a protein that can bind to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor; or (c) a nucleotide sequence which hybridizes with DNA having the nucleotide sequence listed as SEQ. ID. No.1 of the Sequence Listing under stringent conditions and encodes a protein that can bind to an antibody or its fragments that are active to induce granulocyte colony-stimulating factor.

The invention further provides a gene having: (a) the nucleotide sequence listed as SEQ. ID. No.3 of the Sequence Listing; (b) a nucleotide sequence which is the nucleotide

sequence listed as SEQ. ID. No.3 of the Sequence Listing with one or more nucleotide deletions, substitutions, additions or insertions and which encodes a protein that can bind to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor; or (c) a nucleotide sequence which hybridizes with DNA having the nucleotide sequence listed as SEQ. ID. No.3 of the Sequence Listing under stringent conditions and encodes a protein that can bind to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor.

The antibody that is active to induce granulocyte colony-stimulating factor mentioned above is, for example, the monoclonal antibody produced by a hybridoma of the cell line deposited as FERM BP-6103.

According to the invention, the gene is a gene derived from a mouse or human.

The invention further provides a DNA fragment containing: (1) the nucleotide sequence from position 519 to position 736, the nucleotide sequence from position 666 to position 689, the nucleotide sequence from position 381 to position 403 or the nucleotide sequence from position 709 to position 727 of the nucleotide sequence listed as SEQ. ID. No.1 of the Sequence Listing; (2) a nucleotide sequence which is any of the nucleotide sequences of (1) above with one or more nucleotide deletions, substitutions, additions or insertions; or (3) a nucleotide sequence which has at least 80% homology with any of the nucleotide sequences of (1) above.

The invention further provides a gene containing: (1) the nucleotide sequence from position 519 to position 736, the nucleotide sequence from position 666 to position 689, the nucleotide sequence from position 381 to position 403  
5 or the nucleotide sequence from position 709 to position 727 of the nucleotide sequence listed as SEQ. ID. No.1 of the Sequence Listing; (2) a nucleotide sequence which is any of the nucleotide sequences of (1) above with one or more nucleotide deletions, substitutions, additions or  
10 insertions; or (3) a nucleotide sequence which has at least 80% homology with any of the nucleotide sequences of (1) above; and encoding a protein that can bind to an antibody or its fragments that are active to induce granulocyte colony-stimulating factor.

15 The invention further provides any of the following proteins: (a) a protein having the amino acid sequence listed as SEQ. ID. No.2 of the Sequence Listing; (b) a protein having the amino acid sequence listed as SEQ. ID. No.2 of the Sequence Listing with one or more amino acid  
20 deletions, substitutions, additions or insertions and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor; (c) a protein having at least 50% homology with the amino acid sequence listed as SEQ. ID. No.2 and also binding to an antibody or  
25 its fragment that is active to induce granulocyte colony-stimulating factor; or (d) a protein that is encoded by DNA which hybridizes with DNA having the nucleotide sequence listed as SEQ. ID. No.1 of the Sequence Listing under



stringent conditions and that binds to an antibody or its fragments that are active to induce granulocyte colony-stimulating factor.

The invention further provides any of the following  
5 proteins: (a) a protein having the amino acid sequence listed as SEQ. ID. No.4 of the Sequence Listing; (b) a protein having the amino acid sequence listed as SEQ. ID. No.4 of the Sequence Listing with one or more amino acid deletions, substitutions, additions or insertions and also  
10 binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor; (c) a protein having at least 50% homology with the amino acid sequence listed as SEQ. ID. No.4 and also binding to an antibody or its fragment that is active to induce granulocyte colony-  
15 stimulating factor; or (d) a protein that is encoded by DNA which hybridizes with DNA having the nucleotide sequence listed as SEQ. ID. No.3 of the Sequence Listing under stringent conditions and that binds to an antibody or its fragment that is active to induce granulocyte colony-  
20 stimulating factor.

The antibody that is active to induce granulocyte colony-stimulating factor mentioned above is, for example, the monoclonal antibody produced by a hybridoma of the cell line deposited as FERM BP-6103.

25 According to the invention, the protein is preferably a protein derived from mammals and most preferably from a mouse or human.

The invention further provides a protein comprising

any of the followings: (1) the amino acid sequence from residues 1 to 91, the amino acid sequence from residues 50 to 146, the amino acid sequence from residues 1 to 78, the amino acid sequence from residues 200 to 241, the amino acid sequence from residues 172 to 241, the amino acid sequence from residues 103 to 150 or the amino acid sequence from residues 169 to 241 of the amino acid sequence listed as SEQ. ID. No.2 of the Sequence Listing; (2) an amino acid sequence which is any of the amino acid sequences of (1) above with one or more amino acid deletions, substitutions, additions or insertions; or (3) an amino acid sequence having at least 70% homology with any of the amino acid sequences of (1) above.

The invention further provides a protein comprising any of the followings: (1) the amino acid sequence from residues 1 to 91, the amino acid sequence from residues 50 to 146, the amino acid sequence from residues 1 to 78, the amino acid sequence from residues 200 to 241, the amino acid sequence from residues 172 to 241, the amino acid sequence from residues 103 to 150 or the amino acid sequence from residues 169 to 241 of the amino acid sequence listed as SEQ. ID. No.2 of the Sequence Listing; (2) an amino acid sequence which is any of the amino acid sequences of (1) above with one or more amino acid deletions, substitutions, additions or insertions; or (3) an amino acid sequence having at least 70% homology with any of the amino acid sequences of (1) above, and also binding to an antibody or its fragments that are active to

induce granulocyte colony-stimulating factor.

The invention further provides an antibody or fragment thereof, for the aforementioned protein of the invention. The antibody is preferably a monoclonal  
5 antibody, and most preferably a human-type monoclonal antibody or human monoclonal antibody.

The invention further provides a recombinant vector containing a gene or a DNA fragment according to the invention.

10 The invention further provides a transformant comprising a recombinant vector containing a gene or a DNA fragment according to the invention.

The invention further provides a receptor for a substance that can induce production of granulocyte colony-  
15 stimulating factor, comprising a protein according to the invention.

The invention further provides a method employing a protein of the invention for screening of a useful substance (for example, an agonist or antagonist for the  
20 protein), the substance obtained by the screening method and a useful substance that can bind to a receptor (for example, an agonist or antagonist for the receptor).

The invention further provides a pharmaceutical compounds comprising a gene, a DNA fragment, a protein (or  
25 a protein fragment), an antibody (or an antibody fragment), a receptor or a substance according to the invention (particularly a pharmaceutical compounds for diagnosis, prevention or treatment of infectious diseases or

neutropenia), and a treatment method employing it.

#### DETAILED DESCRIPTION OF THE INVENTION

Embodiments and working methods of the invention will  
5 now be explained in detail.

Prior to the present invention, the present inventors  
succeeded in obtaining antibodies by direct immunization of  
macrophages and isolating G-CSF inducing antibodies among  
the obtained antibodies (Japanese Patent Application HEI No.  
10 9-266591, all of the content of which is incorporated in  
the present specification by reference). The gene of the  
present invention was isolated by using these antibodies as  
a probe for screening of a cDNA library derived from mouse  
macrophages, and the protein encoded by the gene of the  
15 invention is characterized by the ability to bind to the  
antibodies or the fragments thereof which have granulocyte  
colony-stimulating factor-inducing activity.

(Antibodies or fragments thereof which have granulocyte  
20 colony-stimulating factor-inducing activity)

First, an explanation will be provided regarding the  
method of obtaining the "antibodies or fragments which have  
granulocyte colony-stimulating factor-inducing activity"  
according to the invention. (Also referred to as  
25 "antibodies used for the invention" hereinafter throughout  
the present specification.)

The present inventors administered a mouse macrophage  
cell line as an immunogen to MRL/lpr mice (autoimmune mice),

and isolated monoclonal antibodies. Then, the obtained monoclonal antibodies were applied to the mouse macrophage cell line and examined the effect of the antibodies to the macrophage cells. As a result, it was discovered that one  
5 of the obtained antibodies had the character of causing production of G-CSF by the mouse macrophage cell line in a concentration-dependent manner. (The hybridoma cell line that produces the antibody has been deposited as FERM BP-6103.)

10 Throughout the present specification, the term "monoclonal antibodies" will refer to monoclonal antibodies with reactivity to the macrophage cell line, and specifically refer to monoclonal antibodies with a function of causing production of G-CSF.

15 The antibodies used for the invention had the character to bind substantially to the macrophage cell line. The antibodies used for the invention include any polyclonal antibody or monoclonal antibody having this characters. The "monoclonal antibodies" include monoclonal  
20 antibodies belonging to all the immunoglobulin classes IgG, IgM, IgA, IgD and IgE, and are preferably monoclonal antibodies of the immunoglobulin classes IgG and IgM.

The macrophage cell line may be prepared from naturally occurring leukemia cells or it may be prepared by  
25 transformation with leukemia virus.

The antibodies used for the invention may be obtained according to a common procedure (for example, the method described in "Zoku Seikagaku Jikken Koza 5, Men'eki



antibody-producing cells are collected from the immunosensitized animals after about 3-5 days from the final immunization.

The monoclonal antibodies used for the invention include the monoclonal antibodies produced by a hybridoma of "FERM BP-6103" (3-4H7 antibodies), their fragments and antibodies having essentially the same characters. "3-4H7 antibody" has the ability to induce G-CSF production by the cells.

10 The hybridoma that produces the monoclonal antibodies used for the invention may be prepared by a commonly known method. As an example of a commonly known method for preparation of a monoclonal antibody-secreting hybridoma, there may be mentioned in the methods of Koehler and  
15 Milstein (Nature, Vol. 256, pp.495-497, 1975) or its modified methods. Here, the monoclonal antibodies are prepared by culturing fused cells (a hybridoma) that are obtained by fusing antibody-producing cells from a spleen, lymph node, bone marrow or tonsil, preferably a spleen,  
20 taken from an animal immunosensitized as described above, with myeloma cells from a mammalian animal such as a mouse, rat, guinea pig, hamster, rabbit or human, preferably of the same species, and more preferably from a mouse, rat, or human. The culturing may be carried out *in vitro*, or *in*  
25 *vivo* in the ascites fluid of a mammalian animal such as a mouse, rat, guinea pig, hamster or rabbit, preferably a mouse or rat, and more preferably a mouse, and the antibodies may be obtained from each culture supernatants

or from the ascites fluid of the mammalian animal.

As examples of myeloma cell lines to be used for the cell fusion, there may be the mouse-derived myelomas "P3/X63-AG8", "P3/NSI/1-Ag4-1", "P3/X63-Ag8.U1", "SP2/0-Ag14", "PAI", "FO" or "BW5147", the rat-derived myeloma "210RCY3-Ag1.2.3" and the human-derived myelomas "U-266AR1", "GM1500-6TG-A1-2", "UC729-6", "CEM-AGR", "D1R11" and "CEM-T15".

The screening of fused cell clones that produce the monoclonal antibodies and used for the invention may be accomplished by measuring the antigen reactivity of the culture supernatants from the wells exhibiting cell growth in a microtiter plate using an enzyme-immunological method such as flow cytometry, RIA or ELISA.

As examples of basic media, there may be low calcium media such as Ham'F12 medium, MCDB153 medium or low calcium MEM medium, and high calcium media such as MCDB104 medium, MEM medium, D-MEM medium, RPMI1640 medium, ASF104 medium or RD medium. Serum, hormone cytokines and/or various organic or inorganic substances may be added to the basic medium depending on the purpose. Isolation and purification of the monoclonal antibodies from the culture supernatant or ascites fluid can be accomplished by various method such as saturated ammonium sulfate method, euglobulin precipitation, caproic acid method, caprylic acid method, ion-exchange chromatography (DEAE, DE52, etc.), affinity column chromatography with an anti-immunoglobulin column or Protein A or Protein G column, or hydrophobic chromatography.



The monoclonal antibodies used for the invention may also be obtained by any other method, without being restricted to the production method described above. Ordinary "monoclonal antibodies" have sugar chains with  
5 different structures depending on the types of a mammalian animal which has been immunosensitized, and the "monoclonal antibodies" used for the invention are not limited by structural differences in these sugar chains and include any monoclonal antibody derived from mammalian animals.  
10 The "monoclonal antibodies" used for the invention also include monoclonal antibodies produced by phage display, as well as human-type monoclonal antibodies obtained using transgenic mice created by genetic engineering that produce human-type antibodies by incorporation of human  
15 immunoglobulin genes, for example, chimeric monoclonal antibodies obtained by using gene recombination techniques to recombine the constant region (Fc region) of a mammalian animal-derived monoclonal antibody with the Fc region of a human monoclonal antibody, and humanized monoclonal  
20 antibodies obtained by recombining the complementarity-determining region (CDR) that can directly bind in a complementary manner with the antigen, with the corresponding region of a human monoclonal antibody.

According to the invention, an "antibody fragment"  
25 may also be used, where "antibody fragment" means an antibody fragment including at least one variable region, synonymous with the "antibody portion" mentioned in Japanese Patent Application HEI No. 9-266591. Specifically,

it refers to the Fv, F(ab')<sub>2</sub>, Fab' or Fab fragments. Here, "F(ab')<sub>2</sub>" and "Fab'" refer to antibody fragments produced by treating an immunoglobulin (monoclonal antibody) with a protease such as pepsin or papain, and they are obtained  
5 by digestion before and after the sulfide bonds present between the two H chains at the hinge region. For example, treatment of IgG with papain cleaves it upstream from the disulfide bonds present between the two H chains at the hinge region, producing two homologous antibody fragments  
10 each comprising an L chain composed of a VL (L chain variable region) and CL (L chain constant region) and an H chain fragment composed of a VH (H chain variable region) and CH<sub>γ</sub>1 (γ1 region of the H chain constant region) which are bonded by sulfide bonds at the C-terminal region.  
15 These two homologous antibody fragments are each designated as Fab'. Treatment of IgG with pepsin cleaves it downstream from the disulfide bonds present between the two H chains at the hinge region, producing an antibody fragment which is slightly larger than just the aforementioned two  
20 Fab' fragments connected at the hinge region. This antibody fragment is designated as F(ab')<sub>2</sub>.

The protein encoded by the gene of the invention is characterized by binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating  
25 factor, as explained in detail above. The term "binding" as used throughout the present specification means ordinary binding between a protein and antibody, and it may be measured using common immunological analysis techniques

(for example, immunoprecipitation, ELISA, immunoblotting, etc.)

(Gene of the invention)

5           The present invention provides a gene which encodes the protein having the amino acid sequence listed as SEQ. ID. No.1 of the Sequence Listing or a protein which is homologous thereto. The invention also provides a gene having the nucleotide sequence listed as SEQ. ID. No.1 of  
10 the Sequence Listing or a nucleotide sequence which is homologous thereto.

          There are no particular restrictions on the type of the gene of the invention, and it may be naturally occurring DNA, recombinant DNA or chemically synthesized  
15 DNA, or even a genomic DNA clone or a cDNA clone.

          The gene of the invention will typically have the nucleotide sequence listed as SEQ. ID. No.1 of the Sequence Listing, but this is only the sequence of a clone (MMR19) obtained in the following examples to represent an  
20 embodiment of the invention. Those skilled in the art are well aware that natural genes have a small number of variations depending on the breeding conditions of the biological species that produce it, and on the ecosystem, or on the presence of highly similar isozymes.

25 Consequently, the "gene" of the invention is not limited to the gene having the nucleotide sequence listed as SEQ. ID. No.1 of the Sequence Listing but also includes all genes encoding proteins having the characters described in the

present specification.

In particular, disclosure by the present specification of the amino acid sequence for the protein of the invention and the DNA sequence encoding the protein easily  
5 allows to isolate a gene encoding a protein with similar physiological activity from another biological species by utilizing basic genetic engineering techniques such as hybridization or PCR. Genes obtained in this manner are also within the scope of the present invention.

10 There are no particular restrictions for the hybridization conditions used for screening of a homologous gene, and they may be appropriately selected by a person skilled in the art depending on the degree of homology between the target homologous gene and probe, although stringent  
15 conditions are generally preferred, and for example, the hybridization conditions may be 6x SSC [0.9 M NaCl, 0.09 M sodium citrate (pH 7.0)], 5 x Denhardt's solution [1 g ficoll, 1 g polyvinyl pyrrolidone, 1 g BSA in 1000 mL], 0.5% SDS, 25°C-68°C (for example, 37°C, 42°C or 68°C), or 0-  
20 50% formamide, 6x SSC, 0.5% SDS, 25-68°C (for example, 37°C, 42°C or 68°C). It is well known to those skilled in the art that appropriate setting of the hybridization conditions including the formamide concentration, Denhardt's solution concentration, salt concentration, temperature, etc. will  
25 allow to clone DNA containing a nucleotide sequence with a given degree of homology or greater, and all homologous genes cloned in this manner are within the scope of the present invention.

A homologous gene cloned by hybridization in this manner has at least 70%, preferably at least 80%, more preferably at least 90%, even more preferably at least 95% and most preferably at least 98% homology with respect to  
5 the nucleotide sequence listed as SEQ. ID. No.1 of the Sequence Listing.

(Protein of the invention)

The present invention provides the protein having the  
10 amino acid sequence listed as SEQ. ID. No.1 of the Sequence Listing or a protein homologous thereto.

The protein having the amino acid sequence listed as SEQ. ID. No.1 of the Sequence Listing according to the invention may be obtained by incorporating the gene  
15 encoding therefor into an appropriate expression vector, transforming this vectors to an appropriate host and expressing the recombinant protein. However, the source and preparation method are not restricted so long as the protein of the invention has the characters described in  
20 the present specification, and it may be a naturally produced protein, a protein expressed from recombinant DNA by a genetic engineering method or a protein chemically synthesized.

The protein of the invention will typically have the  
25 sequence of 241 amino acids listed as SEQ. ID. No.1 of the Sequence Listing. However, those skilled in the art are well aware that natural proteins include variations of one or more amino acids due to gene variation depending on the

breeding conditions of biological species that produce it, on the ecosystem, the presence of highly similar isozymes. The term "amino acid variation" as used here means one or more amino acid substitutions, deletions, insertions and/or  
5 additions. The "protein" of the invention has the amino acid sequence listed as SEQ. ID. No.1 based on deduction from the nucleotide sequence of the cloned gene, but it is not limited only to proteins with that sequence and is intended to include all homologous proteins that have the  
10 characteristics described in the present specification. The homology is at least 50%, preferably at least 60%, more preferably at least 70%, more preferably at least 80%, even more preferably at least 90%, yet more preferably at least 95% and most preferably at least 98%.

15 Generally speaking, introduction of a substitution between amino acids with the same nature (for example, a substitution between two hydrophobic amino acids, a substitution between two hydrophilic amino acids, a substitution between two acidic amino acids or a  
20 substitution between two basic amino acids) will usually tend to give a varied protein with the same characters as the original protein. Those skilled in the art are familiar with methods of preparing recombinant proteins with desired characters using gene recombinant techniques,  
25 and such varied proteins are also within the scope of the invention.

The following examples in the present specification describe cloning of mouse macrophage-derived cDNA as an

embodiment of the invention. The use of the amino acid sequence of the protein disclosed in the present specification or the sequence of the (mouse-derived) gene coding therefor, or a portion thereof, for isolation of a  
5 gene encoding a protein from another source but having similar physiological activity using gene engineering techniques such as hybridization or PCR, is within the scope of commonly accepted knowledge to a person skilled in the art, and proteins encoded by such isolated genes are  
10 also within the scope of the present invention.

(Human-type gene and protein)

For example, the following method may be mentioned as an example of obtaining a human-derived homologue of the  
15 gene and protein of the invention.

The total RNA is extracted from a human macrophage cell line (THP-1, U937, HL-60) by guanidium thiocyanate-phenol-chloroform single-step extraction (Laboratory  
Manuals of Genetic Engineering, 3rd Edition, pp.83-84,  
20 1996), and purified using an oligo(dT) cellulose column, to obtain poly(A)<sup>+</sup> RNA. Reverse transcriptase (MMLV-RTase) and DNA polymerase are used to synthesize double-stranded cDNA. The double-stranded cDNA is used to construct a cDNA library using a  $\lambda$ ZAPII phage vector by the method of  
25 Gubler-Hoffmann (Gubler, U. and Hoffmann, B.J.: Gene, 25:263-269, 1983). A probe is then prepared by amplifying a DNA sequence using a primer DNA that can amplify a sequence in the region of the nucleotide sequence (SEQ. ID.

No.1) of the mouse cDNA (MMR19 clone) disclosed in the present specification, having high homology with the human sequence (for example, the region from position 172 to position 241 of SEQ. ID. No.1 which has been found to have 5 91% homology with the human sequence) and the template DNA from the human macrophage cell cDNA library. Or region (for example, the region from position 172 to position 241 of SEQ. ID. No.1) is used directly as the probe for screening the cDNA encoding the entire length of the target 10 protein from the human macrophage cell cDNA library. The cDNA nucleotide sequence is analyzed by the Primer Walking method. The cDNA confirmed to encode the entire length of the target protein is introduced into a baculovirus to express a protein, which can be purified with an affinity 15 column to obtain the human-type homologous protein.

As explained above, the present invention relates to the gene or protein having the nucleotide sequence listed as SEQ. ID. No.1 or the amino acid sequence listed as SEQ. ID. No.2, and to genes and proteins which are homologous 20 thereto. As a result of a search to determine whether or not sequences homologous to the nucleotide sequence listed as SEQ. ID. No.1 and the amino acid sequence listed as SEQ. ID. No.2 provided by the invention are present in other organisms, it was confirmed that human ESTs (expressed 25 sequence tags) include sequences having high homology with the gene of the invention (see Example 3 below). It will therefore be readily appreciated that the human-derived homologous gene can also be isolated by screening of a



human-derived gene library (cDNA library, etc.) using the human-derived ESTs having high homology with the nucleotide sequence of the invention as the probe.

As described above, a database search revealed that  
5 portions (i.e. DNA fragments) of the nucleotide sequence listed as SEQ. ID. No.1 according to the invention are conserved, having high homology with the human sequence. Such DNA fragments are useful as probes for screening of the human-derived homologous gene, and therefore constitute  
10 one aspect of the present invention. The DNA fragments include DNA fragments containing the nucleotide sequence from position 519 to position 736, the nucleotide sequence from position 666 to position 689, the nucleotide sequence from position 381 to position 403 or the nucleotide  
15 sequence from position 709 to position 727 of the nucleotide sequence listed as SEQ. ID. No.1 of the Sequence Listing, while DNA fragments containing a nucleotide sequence which is any of these nucleotide sequences with one or more nucleotide deletions, substitutions, additions  
20 or insertions or a nucleotide sequence which has at least 80%, preferably at least 85%, more preferably at least 90%, even more preferably at least 95% and most preferably at least 98% homology with any of these nucleotide sequences, are also within the scope of the invention.

25 A database search also revealed that a portion of the amino acid sequence listed as SEQ. ID. No.2 according to the invention is conserved with high homology in the human sequence. Protein fragments comprising portions of the

protein of the invention are also useful as reagents for analysis and isolation of antibodies with G-CSF inducing activity, as is the protein of the invention, and also have potential utility as a drug like the protein of the  
5 invention, and thus constitute an aspect of the invention.

The mentioned proteins include the amino acid sequences from residues 1 to 91, 50 to 146, 1 to 78, 200 to 241, 172 to 241, 103 to 150, and 169 to 241 of the amino acid sequence listed as SEQ. ID. No.2 of the Sequence  
10 Listing, while proteins containing an amino acid sequence which is any of these amino acid sequences with one or more amino acid deletions, substitutions, additions or insertions or an amino acid sequence having at least 70%, preferably at least 80%, more preferably at least 85%, even  
15 more preferably at least 90%, even more preferably at least 95% and most preferably at least 98% homology with any of these amino acid sequences, are also within the scope of the invention.

The present inventors have determined the nucleotide  
20 sequence of the human-type antigen gene by a method similar to the one described above (see Example 4 below). Consequently, the present invention provides a gene having the nucleotide sequence listed as SEQ. ID. No.3 of the Sequence Listing or having a nucleotide sequence which is  
25 homologous thereto. The invention also provides a protein having the amino acid sequence listed as SEQ. ID. No.4 of the Sequence Listing or a protein which is homologous thereto. Here, "homologous" means that the scope of the

invention is not limited to the gene having the nucleotide sequence listed as SEQ. ID. No.3 or the protein having the amino acid sequence listed as SEQ. ID. No.4, as was explained in the part of "Gene of the invention" and  
5 "Protein of the invention" described above.

(Antibodies of the invention)

The invention also provides antibodies against the above-mentioned protein of the invention (also referred to  
10 hereinafter in the present specification as "monoclonal antibodies of the invention"). An example of the antibodies of the invention and a method for obtaining them is explained in detail.

The antibodies of the invention may be either poly-  
15 clonal antibodies or monoclonal antibodies, and in the case of monoclonal antibodies they may be chimeric antibodies. Mouse/human chimeric antibodies are particularly preferred. The "monoclonal antibodies" include monoclonal antibodies belonging to all the immunoglobulin classes such as IgG,  
20 IgM, IgA, IgD and IgE, and are preferably monoclonal antibodies of the immunoglobulin classes IgG and IgM.

The protein of the invention used as the antigen may be obtained by incorporating a gene encoding it into an appropriate expression vector, transforming the  
25 incorporated vector to an appropriate host and expressing the recombinant protein. The immunogen used may be, for example, an actual macrophage cell line or the membrane fraction of a macrophage cell line.

The antibodies of the invention, such as polyclonal antibodies (antiserum) or monoclonal antibodies may be obtained according to a common procedure (for example, the method described in "Zoku Seikagaku Jikken Koza 5, Men'eki  
5 Seikagaku Kenkyuuhou" [Method of Biochemical Experiments V - Immunobiochemistry Research Methods], ed. by The Japanese Biochemical Society: published by Tokyo Kagaku Dojin).

Specifically, a mammalian animal, preferably a mouse, rat, hamster, guinea pig, rabbit, dog, cat, pig, goat,  
10 horse or cow, and more preferably a mouse, rat, hamster, guinea pig or rabbit, is immunized with the antigen in combination with Freund's adjuvant if necessary.

Polyclonal antibodies may be collected from serum obtained from the immunosensitized animal. Monoclonal antibodies  
15 may be produced from a hybridoma made of fusions between antibody-producing cells obtained from the immunosensitized animal and a myeloma cell line (myeloma cells) which has no ability to produce antibodies. The hybridoma was cloned the hybridoma and selected clones that produce monoclonal  
20 antibodies exhibiting specific affinity toward the antigen used to immunize the mammalian animal.

Specifically, the monoclonal antibodies may be produced in the following manner. The protein of the invention or cells expressing the protein of the invention  
25 are used as the immunogen together with Freund's adjuvant if necessary. For immunosensitization, a mouse, rat, hamster, guinea pig or rabbit, and preferably a mouse, rat or hamster (such animals include transgenic animals created

to produce antibodies of other animals, such as human antibody-producing transgenic mice) is used by one or several injections through a subcutaneous, intramuscular, intravenous, foot pad or intraabdominal route, or by  
5 transplantation. Usually 1 to 4 booster immunizations are given every 1 to 14 days from the initial immunization, and 1 to 5 days after the final immunization, and the antibody-producing cells are taken from the immunosensitized mammalian animal.

10 The monoclonal antibodies of the invention may be produced from a hybridoma (fused cells) produced by cell fusion.

The hybridoma producing the monoclonal antibodies can be prepared by a commonly known method. As a commonly  
15 known method, it may be mentioned in the method of Koehler and Milstein (Nature, Vol. 256, pp.495-497, 1975) or methods with modifications of that method. Specifically, the monoclonal antibodies are prepared by culturing fused cells (a hybridoma), which are obtained by fusing antibody-  
20 producing cells from a spleen, lymph nodes, bone marrow or tonsils, preferably a spleen, taken from an animal immunosensitized in the manner described above, with myeloma cells from a mammalian animal such as a mouse, rat, guinea pig, hamster, rabbit or human, and preferably from a  
25 mouse, rat, or human.

As examples of myeloma cell lines used for the cell fusion, there may be the mouse-derived myelomas as mentioned above, such as "P3/X63-AG8", "P3/NSI/1-Ag4-1",

"P3/X63-Ag8.U1", "SP2/0-Ag14", "X63,653", "PAI", "FO" or "BW5147", the rat-derived myeloma "210RCY3-Ag1.2.3" and the human-derived myelomas "U-266AR1", "GM1500-6TG-A1-2", "UC729-6", "CEM-AGR", "D1R11" and "CEM-T15".

5           Screening of fused cell clones that produce the monoclonal antibodies used for the invention may be accomplished by culturing the fused cells in a microtiter plate, for example, and by using flow cytometry, RIA, ELISA  
10           alternative the like to measure the antigen reactivity of the culture supernatants from the wells exhibiting growth.

          Production of monoclonal antibodies from the hybridoma may be carried out by culturing the hybridoma *in vitro*, or *in vivo* in the ascites fluid of a mouse, rat, guinea pig, hamster or rabbit, preferably a mouse or rat,  
15           and more preferably a mouse, and isolating the antibodies from the resulting culture supernatant or from the ascites fluid of the mammalian animal. In the case of *in vitro* culture, the hybridoma may be grown, maintained and stored following a condition varied depending on the properties of  
20           the cell line being cultured, the purpose of research and the culturing method. And the culturing used for production of monoclonal antibodies in culture supernatants may be carried out using a known nutrient medium or any nutrient medium derived and prepared from a known basic medium.

25           As examples of basic media, there may be the low calcium media such as Ham'F12 medium, MCDB153 medium or low calcium MEM medium, and high calcium media such as MCDB104 medium, MEM medium, D-MEM medium, RPMI1640 medium, ASF104

medium or RD medium. Serum, hormone cytokines and/or various organic or inorganic substances may also be added to the basic medium, depending on the purpose. Isolation and purification of the monoclonal antibodies can be  
5 accomplished by apply a saturated ammonium sulfate method, euglobulin precipitation, caproic acid method, caprylic acid method, ion-exchange chromatography (DEAE, DE52, etc.), affinity column chromatography with an anti-immunoglobulin column or Protein A or Protein G column, or else subjecting  
10 them to hydrophobic chromatography to subjecting the culture supernatant or ascites fluid.

A "chimeric antibody" according to the invention is a monoclonal antibody created by genetic engineering, and specifically, it refers to a chimeric monoclonal antibody  
15 such as a mouse/human chimeric monoclonal antibody, which is characterized as the monoglobulin gene whose variable region is a mouse immunoglobulin variable region and the constant region is a human immunoglobulin constant region. The human immunoglobulin constant region is characterized  
20 to have an amino acid sequence depending on the isotype IgG, IgM, IgA, IgD or IgE, and the constant region of the recombinant chimeric monoclonal antibody of the invention may be the constant region of a human immunoglobulin belonging to any of the isotypes. It is preferably the  
25 constant region of human IgG. The chimeric monoclonal antibody of the invention may be produced, for example, in the following manner. It will be appreciated without mention, however, that the production method is not limited

to the one described below.

For example, a mouse/human chimeric monoclonal antibody may be prepared with reference to Jikken Igaku (Experimental Medicine) (special issue) I, Vol.6, N.10, 5 1988 and Japanese Examined Patent Publication HEI No. 3-73280. That is, it may be prepared by inserting into one or separate expression vectors the  $C_H$  gene (C gene encoding the H chain constant region), taken from DNA encoding human immunoglobulin, downstream from the active  $V_H$  gene 10 (rearranged VDJ gene encoding the H chain variable region) taken from DNA encoding the monoclonal antibody isolated from the mouse monoclonal antibody-producing hybridoma, and the  $C_L$  gene (C gene encoding the L chain constant region) taken from DNA encoding human immunoglobulin, downstream 15 from the active  $V_L$  gene (rearranged  $V_J$  gene encoding L chain variable region) taken from DNA encoding the monoclonal antibody isolated from the same hybridoma, with each arranged in an expressible manner, transforming host cells with the expression vector and culturing the transformed 20 cells.

Specifically, first a DNA is extracted from the mouse monoclonal antibody-producing hybridoma by a common procedure, and then the DNA is digested with appropriate restriction endonucleases (for example, EcoRI, HindIII, 25 etc.) and the digested fragments are subjected to electrophoresis (for example, using a 0.7% agarose gel) for Southern blotting. The electrophoresed gel is stained with ethidium bromide, for example, and photographed, after the



marker positions are attached, and the gel is washed twice and then immersed for 15 minutes in a 0.15 M HCl solution. It is then immersed for 10 minutes in a 0.4 N NaOH solution with gentle shaking. A common method is used for transfer  
5 the DNA to a filter, recovery of the filter. After 4 hours, the filter is washed twice with 2x SSC. After thoroughly drying the filter, it is baked (75°C, 3 hours). Upon completion of baking, the filter is placed in a 0.1x SSC/0.1% SDS solution and incubated at 65°C for 30 minutes.  
10 It is then immersed in a 3x SSC/0.1% SDS solution. The obtained filter is placed in a plastic bag together with the prehybridization solution, and incubated at 65°C for 3-4 hours.

Next, <sup>32</sup>P-labelled probe DNA and hybridization  
15 solution are added for reaction at 65°C for about 12 hours. After completion of hybridization, the filter is washed with an appropriate salt concentration, reaction temperature and time (for example, 2x SSC, 0.1% SDS solution, room temperature, 10 minutes). The filter is placed in a  
20 plastic bag, a small amount of 2x SSC is added, the bag is sealed, and autoradiography is performed. This Southern blot method allows identification of the rearranged VDJ gene and VJ gene encoding the H chain and L chain of the mouse monoclonal antibody, respectively. The zones  
25 containing the DNA fragments identified by the method described above are fractionated by sucrose density gradient centrifugation and the isolated DNA is incorporated into a phage vector (for example, charon4A,

charon28,  $\lambda$ EMBL3,  $\lambda$ EMBL4, etc.), and then *E. coli* (for example, LE392, NM539, etc.) is transformed with the phage vector, and a genomic library is created. The genomic library is used for plaque hybridization according to the  
5 method of Benton and Davis (Science, Vol.196, pp.180-182(1977)) using the appropriate probe (H chain J gene, L chain ( $\kappa$ )J gene, etc.), and the positive clones containing either the rearranged VDJ gene or VJ gene are obtained. Restriction enzyme maps of the obtained clones are prepared  
10 and the nucleotide sequences are determined to confirm that the obtained genes contain the target rearranged  $V_H$  (VDJ) gene or  $V_L$  (VJ) gene.

Separately, the human  $C_H$  gene and human  $C_L$  gene used for chimerization are isolated. For example, when creating  
15 a chimera with human IgG1, the  $C\gamma 1$  gene as the  $C_H$  gene and the  $C\kappa$  gene as the  $C_L$  gene are isolated. By taking advantage of the high homology between the nucleotide sequences of the mouse immunoglobulin genes and human immunoglobulin genes, these genes can be obtained using as  
20 probes the mouse  $C\gamma 1$  gene and mouse  $C\kappa$  gene, which correspond to the human  $C\gamma 1$  gene and human  $C\kappa$  gene for their isolation from a human genomic library.

Specifically, the 3 kb HindIII-BamHI fragment from clone Ig146 (Proc. Natl. Acad. Sci. USA, Vol.75, pp.4709-  
25 4713(1978)) and the 6.8 kb EcoRI fragment from clone MEP10 (Proc. Natl. Acad. Sci. USA, Vol.78, pp.474-478(1981)) are used as probes to isolate a DNA fragment containing the human  $\kappa$  gene containing the enhancer region, which is

derived from a human  $\lambda$ Charon4A HaeIII-AluI genomic library (Cell, Vol.15, pp.1157-1174(1978)). The human C $\gamma$ 1 gene is isolated, for example, by digesting human embryonic liver cell DNA with HindIII, fractionating by agarose gel  
5 electrophoresis, and inserting the 5.9 kb band in  $\lambda$ 788 and using the aforementioned probes.

The mouse V $_H$  gene and mouse V $_L$  gene and the human C $_H$  gene and human C $_L$  gene obtained in this manner are incorporated into an expression vector, such as pSV2gpt or  
10 pSV2neo by a common procedure, using an appropriate restriction endonuclease and DNA ligase, taking into account the promoter region and enhancer region, so that the human C $_H$  gene is placed downstream from the mouse V $_H$  gene and the human C $_L$  gene is placed downstream from the  
15 mouse V $_L$  gene. Here, the chimeric genes of the mouse V $_H$  gene/human C $_H$  gene and mouse V $_L$  gene/human C $_L$  gene may be arranged simultaneously in the same expression vector, or they may be arranged in separate expression vectors.

The expression vector inserting the chimeric gene  
20 constructed in this manner is then introduced into myeloma cells such as P3X63·Ag8·653 cells or SP210 cells, which do not of themselves produce antibodies, by the protoplast fusion method, DEAE-dextrin method, calcium sulfate method, electroporation or the like. The transformed cells are  
25 selected out by culturing in medium containing a drug corresponding to the drug resistance gene, which is introduced into the expression vector, and the target chimeric monoclonal antibody-producing cells are isolated.

The target chimeric monoclonal antibodies are taken from the culture supernatant of the selected antibody-producing cells.

The "human-type antibodies (CDR-grafted antibodies)" according to the invention are monoclonal antibodies prepared by genetic engineering, and specifically, they are human-type monoclonal antibodies characterized in that all or a portion of the complementarity-determining region of the hypervariable region is the complementarity-determining region of the hypervariable region derived from the mouse monoclonal antibody, the framework region of the variable region is the framework region of the variable region derived from the human immunoglobulin, and the constant region is the human immunoglobulin region.

The complementarity-determining regions are the three regions found in the hypervariable region of the variable region of the antibody, which are the sites of direct complementary binding to the antigen (CDRs: complementarity-determining regions; CDR1, CDR2, CDR3), and the variable framework regions are the four regions lying before and after the three complementarity-determining regions, which are relatively conserved (Framework regions: FR1, FR2, FR3, FR4). Stated differently, this means a monoclonal antibody wherein all of the regions except for all or a portion of the complementarity-determining region of the hypervariable region of the mouse monoclonal antibody are replaced by the corresponding regions of the human immunoglobulin. The constant region derived from the corresponding region of

the human immunoglobulin has the amino acid sequence characteristic of each isotype IgG, IgM, IgA, IgD or IgE, and the constant region of the human-type monoclonal antibodies of the invention may be the constant region of  
5 human immunoglobulin belonging to any isotype, preferably the constant region of human IgG. There are also no restrictions on the framework regions in the variable region derived from the human immunoglobulin.

The human-type monoclonal antibodies of the invention  
10 may be produced, for example, in the following manner, with understanding that there is no limitation to this production method. For example, the recombinant human-type monoclonal antibodies derived from mouse monoclonal antibodies may be prepared by genetic engineering with  
15 reference to Japanese Patent Public Inspection HEI No. 4-506458 and Japanese Unexamined Patent Publication SHO No. 62-296890. That is, at least one mouse H chain CDR gene and at least one mouse L chain CDR gene pairing to the mouse H chain CDR gene are isolated from the mouse monoclonal  
20 antibody-producing hybridoma, and a human H chain gene encoding the entire region other than the human H chain CDR corresponding to the mouse H chain CDR and the human L chain encoding the entire region other than the human L chain CDR corresponding to the mouse L chain CDR are  
25 isolated from the human immunoglobulin gene.

The isolated mouse H chain CDR gene and human H chain gene are introduced in an expressible manner into an appropriate expression vector, and likewise the mouse L

chain CDR gene and the human L chain gene are introduced in an expressible manner into another appropriate expression vector. Alternatively, the mouse H chain CDR gene/human H chain gene and the mouse L chain CDR gene/human L chain gene may be introduced in an expressible manner into the same expression vector. By transforming host cells with the expression vector prepared in this manner, it is possible to obtain human-type monoclonal antibody-producing transformants, and by culturing these transformants, it is possible to obtain the target human-type monoclonal antibodies from the culture supernatant.

A "human antibody" according to the invention is an immunoglobulin wherein all of the regions of the immunoglobulin including the H chain variable region and H chain constant region and the L chain variable region and L chain constant region are derived from a gene encoding a human immunoglobulin. Human antibodies may be produced by the same method used to prepare polyclonal antibodies or monoclonal antibodies described above. For example, human antibodies are made by immunosensitization of a transgenic animal, which was created by incorporating at least a human immunoglobulin gene into the gene locus of a non-human mammalian animal, such as a mouse, according to a common procedure. For example, a human antibody-producing transgenic mouse can be created according to the procedure described in Nature Genetics, Vol.7, pp.13-21, 1994; Japanese Patent Public Inspection HEI No. 4-504365; International Patent Disclosure WO94/25585; Nikkei Science,

No.6, pp.40-50, 1995; Nature, Vol.368, pp.856-859, 1994; or Japanese Patent Public Inspection HEI No. 6-500233.

"Antibody portion" according to the invention means an antibody fragment containing at least one variable  
5 region, and refers to a partial region of an antibody, preferably a monoclonal antibody, according to the invention mentioned above; specifically, it refers to the Fv, F(ab')<sub>2</sub>, Fab' or Fab fragments. Here, "F(ab')<sub>2</sub>" and "Fab'" refer to antibody fragments produced by treating an  
10 immunoglobulin (monoclonal antibody) with a protease such as pepsin or papain, and they are obtained by digestion before and after the sulfide bonds present between the two H chains at the hinge region. For example, treatment of IgG with papain cleaves it upstream from the disulfide  
15 bonds present between the two H chains at the hinge region, resulted in producing two homologous antibody fragments, each consisting of an L chain composed of a V<sub>L</sub> (L chain variable region) and C<sub>L</sub> (L chain constant region) and an H chain fragment composed of a V<sub>H</sub> (H chain variable region)  
20 and C<sub>H</sub>γ1 (γ1 region of the H chain constant region), which are bonded by sulfide bonds at the C-terminal region. These two homologous antibody fragments are both designated as Fab'. Treatment of IgG with pepsin cleaves it downstream from the disulfide bonds present between the two  
25 H chains at the hinge region, resulted in producing an antibody fragment which is slightly larger than the aforementioned two Fab' fragments connected at the hinge region. This antibody fragment is designated as F(ab')<sub>2</sub>.

(Recombinant vector and transformant)

The present invention further provides a recombinant vector containing the gene or DNA fragment of the invention.

The recombinant vector may be prepared by linking the  
5 gene of interest to a recombination vector which is readily  
available to those skilled in the art (for example, plasmid  
DNA or the like) by a common procedure. Examples of  
vectors to be used include, but are not limited to,  
pBluescript, pUC18, pUC19 and pBR322, as plasmids derived  
10 from *E. coli*.

An expression vector is particularly useful for the  
purpose of producing a protein of interest. The type of  
expression vector is not particularly restricted so long as  
it has the function of expressing the gene of interest in  
15 host cells, either or both prokaryotic cells and eukaryotic  
cells, to produce the protein of interest. For example,  
pQE-30, pQE-60, pMAL-C2, pMAL-p2 and pSE420 are preferred  
as expression vectors for *E. coli*, pYES2 (*Saccharomyces*)  
and pPIC3.5K, pPIC9K, pAO815 (all of genus *Pichia*) as  
20 expression vectors for yeast, and pBacPAK8/9, pBK283,  
pVL1392 and pBlueBac4.5 as expression vectors for insects.

As an example of a method for incorporating a gene  
fragment of the invention into a vector such as a plasmid,  
there may be the procedure described in "Sambrook, J. et  
25 al., Molecular Cloning, A Laboratory Manual, (Second  
edition), Cold Spring Harbor Laboratory, 1. 53(1989)". A  
commercially available ligation kit (for example, by Takara  
Shuzo) can be conveniently used. The recombinant vector



(recombinant plasmid, for example) obtained in this manner may be introduced into host cells by the method described below.

Introduction of a recombinant vector of the invention  
5 into host cells (transformation or transfection) can be accomplished by a conventionally known protocol, and as examples, there may be the calcium chloride method or calcium chloride/rubidium chloride method, electroporation, electroinjection, chemical treatment with PEG or the like,  
10 a method using a gene gun, etc., as described in "Sambrook, J. et al., Molecular Cloning, A Laboratory Manual, (Second edition), Cold Spring Harbor Laboratory, 1. 74(1989)". Alternatively, the transformation may be accomplished by the method of Cohen et al. [Proc. Natl. Acad. Sci. USA, 69,  
15 2110(1972)], the protoplast method [Mol. Gen. Genet., 168, 111(1979)] or the competent method [J. Mol. Biol., 56, 209(1971)], for example, when the host cells are bacteria (*E. coli*, *Bacillus subtilis*, etc.); by the method of Hinnen et al. [Proc. Natl. Acad. Sci. USA, 75, 1927(1978)] or the  
20 lithium method [J. Bacteriol., 153, 163(1983)], for example, when the host cells are *Saccharomyces cerevisiae*; by the leaf disk method [Science, 227, 129(1985)] or the electroporation method [Nature, 319, 791(1986)], for example, when the host cells are plant cells; by the method  
25 of Graham [Virology, 52, 456(1973)], for example, when the host cells are animal cells; or by the method of Summers et al. [Mol. Cell. Biol., 3, 2156-2165(1983)], for example, when the host cells are insect cells.

There are no particular restrictions on the host cells to be used to create the transformants so long as they can accommodate and be transformed by the recombinant vector of the invention. Various types of cells may be used, such as naturally existing cells or artificially established recombinant cells, as are commonly used in the technical field of the invention. For example, there may be mentioned prokaryotic cells such as bacteria (*Escherichia*, *Bacillus*) and the like, lower eukaryotic cells including monocellular hosts such as yeast (*Saccharomyces*, *Pichia*) and the like, and higher eukaryotic cells such as silkworm cells and the like. The host cells are preferably *E. coli*, yeast or insect cells, with specific examples including *E. coli* (M15, JM109, BL21, etc.), yeast (INVSc1 (*Saccharomyces*), GS115, KM71 (both of *Pichia*), etc.), and insect cells (BmN4, silkworm larva, etc.). Examples of animal cells include mouse-derived, *Xenopus laevis*-derived, rat-derived, hamster-derived, monkey-derived and human-derived cells, or cultured cell lines established from these cells.

When the host cells are bacteria, particularly *E. coli*, they will usually include at least the expression vector, which has the promoter/operator region, an initiation codon, the gene encoding the protein of interest, a termination codon, a terminator and a replicable unit. When the host cells are yeast, plant cells, animal cells or insect cells, they will usually include at least the expression vector and preferably a promoter, initiation

codon, the gene encoding the protein of interest, a termination codon and a terminator. As appropriate, they may also contain, DNA encoding the signal peptide, an enhancer sequence, the non-translated regions at the 5' and 3' ends of the gene of interest, a selection marker region or a replicable unit.

The preferred initiation codon for the vector of the invention is the methionine codon (ATG). Examples of termination codons are the ordinary termination codons (for example, TAG, TGA and TAA).

A replicable unit means DNA with the ability to replicate its entire DNA sequence in the host cells, and this includes natural plasmids, artificially modified plasmids (plasmids prepared from natural plasmids) and synthetic plasmids. As preferred plasmids, there may be plasmids pQE30, pET and pCAL or their artificially modified forms (DNA fragments obtained by treating pQE30, pET or pCAL with an appropriate restriction endonuclease), for *E. coli*, plasmids pYES2 and pPIC9K for yeast or plasmid pBacPAK8/9 for insect cells.

The enhancer sequence and terminator sequence used may be ones commonly used by those skilled in the art, such as the ones derived from SV40. The selection marker may be a common one used following an ordinary method. Examples thereof include resistance genes against antibiotics, such as tetracycline, ampicillin, kanamycin, neomycin, hygromycin, spectinomycin or chloramphenicol.

The expression vector may be prepared by linking the

aforementioned promoter, initiation codon, gene encoding the protein of interest, termination codon and terminator region in a continuous and cyclic manner in the appropriate replicable unit. Here the appropriate DNA fragments (for  
5 example, linkers, other restriction enzyme sites, etc.) may be used following a common procedure, such as digestion with a restriction enzyme and ligation using T4DNA ligase, as desired.

10 (Receptor, screening method, novel substance)

The protein encoded by the gene of the invention may possibly act at the entry point of induction and stimulation of G-CSF. (That is, while the present invention is not restricted in any way by the following theory, one  
15 possible explanatory model is that binding of an external ligand to the protein of the invention residing on the surface layer of macrophage cells, transmitting of the resulting signal into the cell, and leading to release of G-CSF by the macrophage.) Consequently, the protein of the  
20 invention could be the receptor working as a granulocyte colony-stimulating factor-inducer or a portion thereof. A "portion of a receptor" would include a subunit of the receptor, possibly modified with a sugar chain or the like. The receptor has the ability to bind (also known as  
25 "affinity") to substances that can allow to induce production of granulocyte colony-stimulating factor, such as monoclonal antibodies produced by the hybridoma deposited as FERM BP-6103 or their fragments, and it may

possibly reside in the cell membrane of cells capable of producing granulocyte colony-stimulating factor, including macrophages. The present invention provides such a receptor.

5

The invention further provides a useful substance screening method by using the protein of the invention. Such a screening method includes measurement of binding between the substance of interest and the protein of the invention, or the receptor, measurement of the effect of the substance of interest via the receptor (for example, production of G-CSF by the macrophages or production of a marker substance from appropriately transformed cells), or comparison between the structure of the substance of interest (for example, its amino acid sequence when the substance of interest is a protein) and the structure of the protein of the invention (for example, its amino acid sequence).

The protein of the invention used for screening is preferably (a) a protein having the amino acid sequence listed as SEQ. ID. No.4 of the Sequence Listing; (b) a protein having the amino acid sequence listed as SEQ. ID. No.4 of the Sequence Listing with one or more amino acid deletions, substitutions, additions or insertions and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor; (c) a protein having at least 50% (preferably at least 60%, more preferably at least 70%, even more preferably at least 80%,

even more preferably at least 90%, especially preferably at least 94% and most preferably at least 98%) homology with the amino acid sequence listed as SEQ. ID. No.4 and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor; or (d) a protein that is encoded by DNA which hybridizes with DNA having the nucleotide sequence listed as SEQ. ID. No.3 of the Sequence Listing under stringent conditions and that binds to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor.

The following is a more specific example of the screening method: A vector is constructed by inserting the G-CSF promoter gene and a gene encoding a marker protein, such as luciferase,  $\beta$ -galactosidase, Green Fluorescent Protein (GFP),  $\beta$ -lactamase or chloramphenicol acetyltransferase (CAT), at downstream thereof, and a drug resistance gene against a drug, such as tetracycline, ampicillin, kanamycin, neomycin, hygromycin or spectinomycin, at further downstream thereof. The vector is introduced into cells (for example, a macrophage cell line, and preferably a human-derived macrophage cell line) bearing receptors, which contains the protein of the invention. The obtained cells are treated with a drug-containing medium, and colony-forming cells are selected. Clones expressing the marker protein upon induction are then selected. It is then confirmed that expression of the marker protein reflects actual expression of G-CSF mRNA. The transformed cell line obtained in this manner is treated with various

substances, followed by screening for substances that have induced expression of the marker protein.

A useful substance obtained by screening is (a) a substance which can bind to the receptor, and as a result  
5 of its binding to the receptor, it can elicit a structural change in the receptor, transmit signals into the cell via the receptor, and induce production of granulocyte colony-stimulating factor (also known as an "agonist" or "agent");  
(b) a substance which can bind to the receptor, and as a  
10 result of its binding to the receptor, it can inhibit the binding of the receptor to the substances that can induce production of granulocyte colony-stimulating factor, but it in itself does not induce production of granulocyte colony-stimulating factor (also known as an "antagonist" or  
15 "blocker"); or (c) a substance which can bind to the receptor, and as a result of its binding to the receptor, it can inhibit the binding of the receptor to the substances that can induce production of granulocyte colony-stimulating factor, but it in itself blocks  
20 production of granulocyte colony-stimulating factor (also known as an "inverse agonist" or "reagent").

Such substances are novel. Thus, the present invention also provides (a) a substance which can bind to  
25 the receptor, and as a result of its binding to the receptor, it can induce a change in the receptor, transmit signals into the cell via the receptor, and induce production of granulocyte colony-stimulating factor; (b) a

substance which can bind to the receptor, and as a result of its binding to the receptor, it can inhibit the binding of the receptor to the substances that can induce production of granulocyte colony-stimulating factor, but it  
5 in itself does not induce production of granulocyte colony-stimulating factor; and (c) a substance which can bind to the receptor, and as a result of its binding to the receptor, it can inhibit the binding of the receptor to the substances that can induce production of granulocyte  
10 colony-stimulating factor, but it in itself blocks production of granulocyte colony-stimulating factor, which are obtained by the method described above. The invention still further provides a substance which can bind to the receptor, and (a) as a result of its binding to the  
15 receptor, it can induce a change in the receptor, transmit signals into the cell via the receptor, and induce production of granulocyte colony-stimulating factor; (b) as a result of its binding to the receptor, it can inhibit the binding of the receptor to the substances that can induce  
20 production of granulocyte colony-stimulating factor, but it in itself does not induce production of granulocyte colony-stimulating factor; or (c) as a result of its binding to the receptor, it can inhibit the binding of the receptor to the substances that can induce production of granulocyte  
25 colony-stimulating factor, but it in itself inhibits production of granulocyte colony-stimulating factor. Such substances will hereinafter be referred to as "substances of the invention".



Examples of substances of the invention include the antibody of the invention, its fragments and other low molecular compounds, among which there are those with the effect of inducing production of granulocyte colony-  
5 stimulating factor, those with the effect of inhibition of receptor binding to the substances that can induce production of granulocyte colony-stimulating factor, and those with the effect of inhibition of receptor binding to the substances that can induce production of granulocyte  
10 colony-stimulating factor, while those also inhibit production of granulocyte colony-stimulating factor.

When the substance of interest is an antibody, the binding to the receptor (or binding inhibition) can be measured by a method, such as, for example, analysis of the  
15 antibody-bound macrophage cells using flow cytometry or ELISA.

Inducing effect (or inhibiting effect) of the production of granulocyte colony-stimulating factor can be determined by the method described in Japanese Unexamined  
20 Patent Publication HEI No. 11-106400. The outline of the process is given below.

The G-CSF promoter gene is inserted between the XhoI and the NcoI site of PicaGene Enhancer Vector 2 (product of Wako Junyaku Kogyo Co., Ltd.), in order to construct vector  
25 Pica G-CSF neo, the luciferase gene is linked downstream therefrom in place of the G-CSF gene itself, and then a neomycin resistance gene cut out from pMC1Neo PolyA at the SalI site downstream from SV40 also. This vector is

introduced into the RAW264.7 cells by electroporation. The obtained cells are treated with medium containing geneticin, and the colony-forming cells are selected. Among the geneticin-resistant clones, clones exhibiting luciferase activity upon induction are further selected. Northern blot analysis using <sup>32</sup>P-labeled mouse G-CSF cDNA as the probe is made to confirm that the luciferase activity reflects actual expression of the G-CSF mRNA. The transformed macrophage cells obtained in this manner are plated in a 96-well microtiter plate at 5 x 10<sup>4</sup> cells per well and cultured at 37°C for 24 hours, and then after treating them with a prepared agonist or antagonist, as need, or the substance of interest is added at concentrations of about 0, 3.75, 7.5, 15, 30 and/or 60 µg/ml. After further culturing at 37°C for 18 hours, the luciferase activity is measured.

(Use of gene of the invention as drug agent)

The gene of the invention may be utilized, for example, for diagnosis, prevention and therapy (gene therapy, etc.) of diseases in which neutrophils, a type of blood leukocyte, are involved (such as neutropenia). The protein, a part of the protein or the peptides thereof, antibody or its fragment, receptor, or the substance of the invention (hereinafter these will sometimes be referred to collectively as "protein, etc. of the invention") can serve as a drug to regulate the number of neutrophils in the blood or bone marrow. That is, the gene and protein, etc. of the invention can be used for treatment of neutropenia

as a side effect of anticancer agents or neutropenia directly following bone marrow transplantation, and for diagnosis, prevention and treatment of anaplastic anemia.

The protein, etc. of the invention may generally be  
5 administered systemically or locally, usually in a parenteral form. Intravenous administration is particularly preferred for the parenteral forms.

The gene of the invention may be administrated systemically or locally in the form of "gene therapy",  
10 wherein the gene is introduced into cells either *in vivo* or *ex vivo*. Introduction of the gene can be accomplished, for example, by the method described in Biomanual UP Series, Idenshi Chiryo no Kiso Gijutsu [Fundamental Techniques for Gene Therapy], Shimada, T., Saito, I., Ozawa, T., ed.:  
15 Yodosha Publishing, 1996. For introduction into cells *ex vivo*, there may be methods employing a retrovirus vector, adenovirus vector, adeno-associated virus (AAV) vector, cationic liposomes, HVJ-liposomes, or the calcium phosphate method, DEAE dextran method, etc. For introduction into  
20 cells *in vivo*, there may be methods employing a retrovirus vector, adenovirus vector, adeno-associated virus (AAV) vector, cationic liposomes or HVJ-liposomes.

The administration dosage will differ depending on age, gender, body weight, symptoms, treatment effect,  
25 administration route, treatment time and substance administrated (types of the protein or gene). But parenteral administration one to several times per day may be given at a dosage in the range of 1  $\mu$ g to 100 g and

preferably in the range of 10  $\mu$ g to 1000 mg for each time for adults. Since the administration dosage will vary depending on the conditions, a dosage below this range will often be sufficient, or a dosage exceeding this range may  
5 be necessary. Injections for parenteral administration according to the invention include sterile aqueous or nonaqueous solutions, suspensions and emulsions. For aqueous and nonaqueous solutions and suspensions, one or more active substances are mixed with at least one inactive  
10 diluting agent. As examples of aqueous diluting agents, there may be distilled water for injection and physiological saline. As examples of nonaqueous diluting agents, there may be propylene glycol, polyethylene glycol, vegetable oils, such as olive oil, and alcohols, such as  
15 ethanol.

Such a composition may also include adjuvants, such as preservatives, humectants, emulsifiers, dispersers and stabilizers (for example, arginine and aspartic acid).

These can be sterilized by passage through a bacteria  
20 capturing filter, mixture with a sterilizer or irradiation. They may also be prepared as sterile solid compositions by lyophilization, for example, and then dissolved in sterile distilled water for injection or another solvent prior to use.

25 Other compositions for parenteral administration include external applications or suppositories and pessaries for enteric administration, which are formulated according to ordinary methods and contain one or more

active substances.

The invention will now be explained in greater detail by way of the following examples, with understanding that  
5 the invention is in no way restricted by these examples.

### Examples

#### Example 1: Cloning of antigen gene recognized by the monoclonal antibody from macrophage cell line

10 (1) Preparation of poly(A)<sup>+</sup> RNA from macrophage cells (RAW264.7)

Guanidium thiocyanate-phenol-chloroform single-step extraction (Laboratory Manuals of Genetic Engineering, 3rd Edition, pp.83-84, 1996) was used to prepare approximately  
15 0.3 mg of total mRNA from 2 x 10<sup>8</sup> mouse macrophage cells (RAW264.7). This was purified using an oligo(dT) cellulose column to obtain 5 µg of poly(A)<sup>+</sup> RNA.

(2) Synthesis of double-stranded cDNA from poly(A)<sup>+</sup>  
20 RNA

A reaction solution (50 µl) containing the poly(A)<sup>+</sup> RNA as obtained above (1) (5 µg), a reverse transcriptase (MMLV-RTase; product of STRATAGENE Corp.; 70 units) and dNTPs (0.6 mM) was incubated at 37°C for 60 minutes to  
25 synthesize of a first strand cDNA. Next, a reaction solution containing the aforementioned reaction solution (45 µl), DNA polymerase (product of STRATAGENE Corp.; 100 units) and dNTPs (0.3 mM) was incubated at 16°C for 150

minutes to synthesize a second strand cDNA, to obtain double-stranded cDNA (8 µg).

### (3) Construction of cDNA library

5       The method of Gubler-Hoffmann (Gubler, U. and Hoffmann, B.J.: Gene, 25:263-269, 1983) was used. PfuDNA polymerase was used to blunt the ends of the double-stranded cDNA as synthesized above (2), and an adapter was linked thereto, by ligation with T4DNA ligase.

10       Specifically, a reaction solution (total: 225 µl) containing the double-stranded cDNA as obtained above (2) (8 µg DNA; 200 µl) and PfuDNA polymerase (5 units) was incubated at 72°C for 30 minutes.

15       The adapter-linked DNA was cleaved at the ends with restriction enzyme XhoI, and the cDNA longer than 0.5 kbp was fractionated with a gel column. This cDNA was incorporated into λZAPII phage vector (STRATAGENE Corp.) with T4DNA ligase following common protocol, and then by ligation into phage particles. Measurement of the phage

20       titer showed that the cDNA library contained  $2 \times 10^6$  independent clones. The obtained phage library was used to infect *E. coli* (XL1-Blue MRF') and was allowed to proliferate to  $3.4 \times 10^9$  pfu/ml.

25       (4) Screening of genes encoding protein that bind to antibodies that show the ability to induce granulocyte colony-stimulating factor

      The cDNA library as constructed above (3) was

subjected to immunoscreening using monoclonal antibodies  
(produced  
by the hybridoma deposited as FERM BP-6103; described in  
Japanese Patent Application HEI No. 9-266591), which shows  
5 the ability to induce granulocyte colony-stimulating factor  
(G-CSF) as the prove. The specific procedure was as  
follows.

*E. coli* (XL1-BlueMRF') infected with the phage cDNA  
library was seeded onto a 150 mm diameter plate. The  
10 plate was incubated at 42°C for 4 hours that allowed the  
formation of the plaques of approximately 0.5 mm diameter.  
Then, microcellulose membranes, which were immersed in  
10 mM of IPTG (isopropylthio- $\beta$ -galactoside) and air-dried,  
were placed on those plates and incubated for 3 hours at  
15 37°C. The nitrocellulose membrane was peeled off, and  
incubated in TBS-T (20 mM Tris-HCl, pH 7.6, 0.1% Tween20)  
containing 5% skim milk for one hour for blocking the  
membranes, while agitating at room temperature. This was  
followed by rinsing of the membrane gently for 2 minutes  
20 with TBS-T (repeated twice), immersing in a buffer at room  
temperature for 15 minutes (once) and washing for 5 minutes  
(twice). The nitrocellulose membrane was incubated for one  
hour at room temperature for reaction with the antibodies  
while agitating in the diluted primary antibody solution  
25 (produced by the hybridoma deposited as FERM BP-6103; 1.6  
 $\mu$ g/ml). The nitrocellulose membrane was then washed in the  
same manner as the previous washing. Alkaline phosphatase-  
labeled secondary antibodies (ZYMED) were diluted to 0.6

µg/ml with TBS containing 1% BSA, and the nitrocellulose membrane was incubated for one hour at room temperature for reaction with the antibodies, while agitating in the diluted secondary antibody solution. The nitrocellulose membrane was again thoroughly washed in the same manner as described above, and finally washed with TBS for 5 minutes. After adding 1 ml of each NBT solution (50 mg/ml of NitroBlue Tetrazolium in 70% dimethylformamide) and BCIP solution (50 mg/ml of 5-bromo-4-chloro-3-indolyl phosphate in dimethylformamide) to a buffer solution containing 100 mM of Tris-HCl (pH 9.5), 100 mM of NaCl and 5 mM of MgCl<sub>2</sub>, the nitrocellulose membrane was immersed therein. Reaction was carried out for 30 minutes in a dark room, and the membrane was washed with water and dried. After drying, the plaques exhibiting positive reaction on the nitrocellulose membrane were collected from the original plate.

Finally, 22 antibody-binding positive clones were obtained from  $7 \times 10^5$  phage in the first screening. The top agarose containing the positive plaques were collected, and amplified. The second, third and fourth screenings were performed from approximately 1000 phages following the same procedure described above, resulting in isolation of 3 positive clones (MMR10, MMR17 and MMR19).

25

(5) Determination of the nucleotide sequence of the gene

The inserts of the 3 positive clones (MMR10, MMR17



and MMR19), as obtained above (4) were cut out from the  
λZAPII phage vectors by *in vivo* excision following a common  
procedure, and subcloned by converting the vector to  
pBluescript SK(-) Phagemid. The subcloned plasmids were  
5 made large quantity in *E. coli* (SOLR), and approximately 20  
μg of the plasmid DNA was obtained. The Primer Walking  
method was used to analyze the nucleotide sequences of  
these DNA.

Based on the results of the nucleotide sequence  
10 analyses, a clone MMR19 was found to have the 840 bp  
nucleotide sequence of the full length cDNA which included  
the open reading frame of the protein. The nucleotide  
sequence of the clone MMR19 is listed as SEQ. ID. No.1 of  
the Sequence Listing.

15

(6) Primary structure of the protein deduced from the  
nucleotide sequence of cDNA clone

The primary structure of the protein (MMR-CAM)  
(listed as SEQ. ID. No.1 and No.2 in the Sequence Listing)  
20 deduced from the nucleotide sequence of the gene analyzed  
(5) consists of 241 amino acid residues, and the molecular  
weight as estimated from the amino acid sequence was  
approximately 27 kDa. MMR-CAM is thought to be a type I  
membrane glycoprotein with one membrane-spanning domain,  
25 which comprises an extracellular portion of 107 amino acids,  
a membrane-spanning portion of 23 amino acids and an  
intracellular portion of 111 amino acids. Homology search  
showed that no molecules were found similar to the protein

of the invention in terms of the structure, suggesting that the protein of the invention does not belong to the existing family. Also, there is a portion with the extensive modifications by the type O sugar chains was present in the extracellular domain. Phosphorylation sites for protein kinase C, tyrosine kinase, etc. are present in the intracellular domain. These sugar chain binding sites and phosphorylation sites are believed to play on very important role in signal transduction.

10

Example 2: Expression of the protein (MMR-CAM) of the invention

The clone (MMR19) obtained in Example 1(4) was inserted into an expression vector ( $\lambda$ ZAPII) following a common procedure and transformed *E. coli* (XL1-Blue), then a transformant cell line was constructed. The transformed *E. coli* cells were cultured, and the culture supernatant was dot blotted and allowed to with the same monoclonal antibodies used as shown in (3), which was produced by hybridoma deposited as FERM BP-6103 as a probe. Following this process, it was confirmed that the culture supernatant contained the protein that bound to the monoclonal antibodies.

25 Example 3: Comparison of the mouse-derived protein with other homologous proteins using database search

A data-search was conducted for human genes homologous to the nucleotide sequence and amino acid

sequence listed as SEQ. ID. No.1 and determined in Example 1 on both the amino acid level and the DNA level databases using (DNA DATA BANK of JAPAN (DDBJ): Dept. of Education, National Institute of Genetics, Center for Information Biology). The results are shown in Tables 1 and 2. These results suggest that the gene of the invention is also conserved in humans with high homology.

Table 1: Homology on amino acid level

Position within amino acid sequence of SEQ. ID. No.1	Matching in human homologue
1 to 91	83/91 (91%)
50 to 146	83/97 (85%)
1 to 78	70/78 (89%)
200 to 241	40/42 (95%)
172 to 241	67/70 (95%)
103 to 150	46/48 (95%)
169 to 241	58/73 (79%)

Table 2: Homology on DNA level

Position within nucleotide sequence of SEQ. ID. No.1	Matching in human homologue
519 to 736	189/218 (86%)
666 to 689	23/24 (95%)
381 to 403	22/23 (95%)
709 to 727	19/19 (100%)

Example 4: Cloning of the human homologue of the antigen gene

Guanidium thiocyanate-phenol-chloroform extraction was used to extract total RNA from human normal brain tissue, and the poly(A)<sup>+</sup> RNA was purified using oligo(dT) cellulose. cDNA was synthesized from the Poly(A)<sup>+</sup> RNA using reverse transcriptase (MMLV-RTase) and DNA polymerase. A sense primer of position 4 to 22 (CCATGTCTGGCTGTCAAGC) and an antisense primer of position 714 to 724 (CCATTTTCTCCAACCTGGGAGC) of the mouse antigen gene (MMR19) sequence were prepared, and these primers and the human normal brain tissue cDNA as the template were used for PCR reaction. As a result, a partial cDNA of the human homologue of the mouse antigen gene (MMR19) was obtained. Next, the 3'RACE method and 5'RACE method were carried out using a specific primer (GSP) for the human homologue partial cDNA and an adapter primer. An antisense primer (GTCAGAAGAGATTTCAGGGTGACC) was prepared from the 3' RACE fragment and a sense primer (AAGCCGTG CGGAGATTGGAGG) from the 5' RACE fragment. As a result of LD-PCR, the full length cDNA of the human homologue including the open reading frame was obtained. The Primer Walking method was used to elucidate the 924 bp nucleotide sequence of the cDNA. The obtained nucleotide sequence is listed as SEQ. ID. No.3 of the Sequence Listing. The nucleotide sequence of the human homologue cDNA (924 bp) showed 84.8% homology (with 712 matching nucleotides out of 924) with the nucleotide sequence of the mouse antigen gene cDNA (840 bp).

The primary structure of the protein deduced from the nucleotide sequence of the obtained gene is listed as SEQ. ID. No.3 and No.4, consisting of 242 amino acids. The deduced amino acid sequence showed 93.8% homology with the mouse form (with 226 matching residues out of 242). This protein is also thought to be a type I membrane glycoprotein with one membrane-spanning domain.

#### EFFECT OF THE INVENTION

10           The gene and the protein encoding by the gene (including fragments of the gene and fragments of the protein), antibody (including fragments thereof), receptor and substance of the invention are novel, and are useful for pharmaceutical purposes.

15           The gene and the protein encoding by the gene (including fragments of the gene and fragments of the protein), antibody (including fragments thereof) and receptor are also useful as analytical reagents for screening of substances (for example, monoclonal antibodies, 20 proteins and other low molecular substances) that have the ability to induce granulocyte colony-stimulating factor.

Fragments of the gene of the invention are also useful as probes for screening of homologous genes derived from other organisms.

# CLAIMS

## 1. A gene encoding:

(a) a protein having the amino acid sequence listed as SEQ. ID. No.2 of the Sequence Listing;

(b) a protein having the amino acid sequence listed as SEQ. ID. No.2 of the Sequence Listing with one or more amino acid deletions, substitutions, additions or insertions and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor; or

(c) a protein having at least 50% homology with the amino acid sequence listed as SEQ. ID. No.2 and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor.

## 2. A gene having:

(a) the nucleotide sequence listed as SEQ. ID. No.1 of the Sequence Listing;

(b) a nucleotide sequence which is the nucleotide sequence listed as SEQ. ID. No.1 of the Sequence Listing with one or more nucleotide deletions, substitutions, additions or insertions and which encodes a protein that can bind to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor; or

(c) a nucleotide sequence which hybridizes with DNA having the nucleotide sequence listed as SEQ. ID. No.1 of the Sequence Listing under stringent conditions and encodes a protein that can bind to an antibody or its fragment that

is active to induce granulocyte colony-stimulating factor.

3. A gene encoding:

(a) a protein having the amino acid sequence listed as SEQ. ID. No.4 of the Sequence Listing;

(b) a protein having the amino acid sequence listed as SEQ. ID. No.4 of the Sequence Listing with one or more amino acid deletions, substitutions, additions or insertions and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor; or

(c) a protein having at least 50% homology with the amino acid sequence listed as SEQ. ID. No.4 and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor.

4. A gene having:

(a) the nucleotide sequence listed as SEQ. ID. No.3 of the Sequence Listing;

(b) a nucleotide sequence which is the nucleotide sequence listed as SEQ. ID. No.3 of the Sequence Listing with one or more nucleotide deletions, substitutions, additions or insertions and which encodes a protein that can bind to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor; or

(c) a nucleotide sequence which hybridizes with DNA having the nucleotide sequence listed as SEQ. ID. No.3 of the Sequence Listing under stringent conditions and encodes a protein that can bind to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor.

5. A gene according to any one of claims 1 to 4, wherein the antibody that is active to induce granulocyte colony-stimulating factor is the monoclonal antibody produced by a hybridoma of the cell line deposited as FERM BP-6103.

6. A gene according to any one of claims 1 to 5, which is a mouse-derived or human-derived gene.

7. A DNA fragment containing:

(1) the nucleotide sequence from position 519 to position 736, the nucleotide sequence from position 666 to position 689, the nucleotide sequence from position 381 to position 403 or the nucleotide sequence from position 709 to position 727 of the nucleotide sequence listed as SEQ. ID. No.1 of the Sequence Listing;

(2) a nucleotide sequence which is any of the nucleotide sequences of (1) with one or more nucleotide deletions, substitutions, additions or insertions; or

(3) a nucleotide sequence which has at least 80% homology with any of the nucleotide sequences of (1).

8. A gene containing:

(1) the nucleotide sequence from position 519 to position 736, the nucleotide sequence from position 666 to position 689, the nucleotide sequence from position 381 to position 403 or the nucleotide sequence from position 709 to position 727 of the nucleotide sequence listed as SEQ. ID. No.1 of the Sequence Listing;

(2) a nucleotide sequence which is any of the nucleotide sequences of (1) with one or more nucleotide deletions, substitutions, additions or insertions; or



(3) a nucleotide sequence which has at least 80% homology with any of the nucleotide sequences of (1), and encoding a protein that can bind to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor.

9. Any of the following proteins:

(a) a protein having the amino acid sequence listed as SEQ. ID. No.2 of the Sequence Listing;

(b) a protein having the amino acid sequence listed as SEQ. ID. No.2 of the Sequence Listing with one or more amino acid deletions, substitutions, additions or insertions and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor;

(c) a protein having at least 50% homology with the amino acid sequence listed as SEQ. ID. No.2 and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor; or

(d) a protein that is encoded by the DNA which hybridizes with DNA having the nucleotide sequence listed as SEQ. ID. No.1 of the Sequence Listing under stringent conditions and that binds to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor.

10. Any of the following proteins:

(a) a protein having the amino acid sequence listed as SEQ. ID. No.4 of the Sequence Listing;

(b) a protein having the amino acid sequence listed

as SEQ. ID. No.4 of the Sequence Listing with one or more amino acid deletions, substitutions, additions or insertions and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor;

(c) a protein having at least 50% homology with the amino acid sequence listed as SEQ. ID. No.4 and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor; or

(d) a protein that is encoded by the DNA which hybridizes with DNA having the nucleotide sequence listed as SEQ. ID. No.3 of the Sequence Listing under stringent conditions and that binds to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor.

11. A protein according to claim 9 or 10, wherein the antibody that is active to induce granulocyte colony-stimulating factor is the monoclonal antibody produced by a hybridoma of the cell line deposited as FERM BP-6103.

12. A protein according to any one of claims 9 to 11, which is a mouse-derived, human-derived or other mammalian-derived protein.

13. A protein comprising any of the followings:

(1) the amino acid sequence from residues 1 to 91, the amino acid sequence from residues 50 to 146, the amino acid sequence from residues 1 to 78, the amino acid sequence from residues 200 to 241, the amino acid sequence from residues 172 to 241, the amino acid sequence from

residues 103 to 150 or the amino acid sequence from residues 169 to 241 of the amino acid sequence listed as SEQ. ID. No.2 of the Sequence Listing;

(2) an amino acid sequence which is any of the amino acid sequences of (1) with one or more amino acid deletions, substitutions, additions or insertions; or

(3) an amino acid sequence having at least 70% homology with any of the amino acid sequences of (1).

14. A protein comprising any of the followings:

(1) the amino acid sequence from residues 1 to 91, the amino acid sequence from residues 50 to 146, the amino acid sequence from residues 1 to 78, the amino acid sequence from residues 200 to 241, the amino acid sequence from residues 172 to 241, the amino acid sequence from residues 103 to 150 or the amino acid sequence from residues 169 to 241 of the amino acid sequence listed as SEQ. ID. No.2 of the Sequence Listing;

(2) an amino acid sequence which is any of the amino acid sequences of (1) with one or more amino acid deletions, substitutions, additions or insertions; or

(3) an amino acid sequence having at least 70% homology with any of the amino acid sequences of (1), and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor.

15. An antibody against a protein according to any one of claims 9 to 14, or a fragment thereof.

16. An antibody or fragment thereof according to claim 15, which is a monoclonal antibody.

17. An antibody or fragment thereof according to claim 16, which is a human-type monoclonal antibody or human monoclonal antibody.

18. A recombinant vector containing a gene or DNA fragment according to any one of claims 1 to 8.

19. A transformant comprising a recombinant vector that contains the gene or the DNA fragment according to any one of claims 1 to 8.

20. A receptor for a substance that can induce production of granulocyte colony-stimulating factor including, such as a monoclonal antibody or its fragment that is produced by a hybridoma of the cell line deposited as FERM BP-6103, and the receptor comprises a protein according to any one of claims 9 to 12 and is present in a cell which can produce granulocyte colony-stimulating factor, such as a macrophage.

21. A screening method for any of the following substances, the method includes measurement of binding between a substance and a protein according to any one of claims 9 to 12 or a receptor according to claim 20, measurement of the effect of the substance via a receptor according to claim 20, or measurement to compare the effect between the structure of a substance and the structure of a protein according to the invention.

(a) a substance which can bind to a receptor according to claim 20, and as a result of its binding to a receptor, it can cause a change in the receptor structure, transmit signals into the cell via the receptor, and induce production of granulocyte colony-stimulating factor;

(b) a substance which can bind to a receptor according to claim 20, and as a result of its binding to a receptor, it can inhibit the binding of the receptor to the substances that can induce production of granulocyte colony-stimulating factor, but it in itself does not induce production of granulocyte colony-stimulating factor; or

(c) a substance which can bind to a receptor according to claim 20, and as a result of its binding to a receptor, it can inhibit the binding of the receptor to the substances that can induce production of granulocyte colony-stimulating factor, but it in itself blocks production of granulocyte colony-stimulating factor.

22. Any of the following substances obtained by a screening method, which is characterized by measurement of binding between a substance and a protein according to any one of claims 9 to 12 or a receptor according to claim 20, or measurement of the effect of the substance via a receptor according to claim 20.

(a) a substance which can bind to a receptor according to claim 20, and as a result of its binding to a receptor, it can cause a change in the receptor structure, transmit signals into the cell via the receptor, and induce production of granulocyte colony-stimulating factor;

(b) a substance which can bind to a receptor according to claim 20, and as a result of its binding to a receptor, it can inhibit the binding of the receptor to the substances that can induce production of granulocyte colony-stimulating factor, but it in itself does not induce

production of granulocyte colony-stimulating factor; or

(c) a substance which can bind to a receptor according to claim 20, and as a result of its binding to a receptor, it can inhibit the binding of the receptor to the substances that can induce production of granulocyte colony-stimulating factor, but it in itself blocks production of granulocyte colony-stimulating factor.

23. Any of the following substances that can bind to a receptor according to claim 20:

(a) a substance which, as a result of its binding to a receptor, it can cause a change in the receptor structure, transmit signals into the cell via the receptor, and induce production of granulocyte colony-stimulating factor;

(b) a substance which, as a result of its binding to a receptor, it can inhibit the binding of the receptor to the substances that can induce production of granulocyte colony-stimulating factor, but it in itself does not induce production of granulocyte colony-stimulating factor; or

(c) a substance which, as a result of its binding to a receptor, it can inhibit the binding of the receptor to the substances that can induce production of granulocyte colony-stimulating factor, but it in itself blocks production of granulocyte colony-stimulating factor.

24. A pharmaceutical composition comprising a gene or DNA fragment according to any one of claims 1 to 8, a protein according to any one of claims 9 to 14, an antibody or its fragment according to any one of claims 15 to 17, a receptor according to claim 20 or a substance according to

claim 22 or 23.

25. A pharmaceutical composition according to claim 23 which is used for diagnosis, prevention or treatment of a disease or condition related to G-CSF, such as infection or neutropenia.

26. A method of diagnosis, prevention or treatment of a disease or a condition related to G-CSF, such as infection or neutropenia, which employs a gene or DNA fragment according to any one of claims 1 to 8, a protein according to any one of claims 9 to 14, an antibody or its fragment according to any one of claims 15 to 17, a receptor according to claim 20 or a substance according to claim 22 or 23.

27. The use of a gene or DNA fragment according to any one of claims 1 to 8, a protein according to any one of claims 9 to 14, an antibody or its fragment according to any one of claims 15 to 17, a receptor according to claim 20 or a substance according to claim 22 or 23, for diagnosis, prevention or treatment of a disease or condition related to G-CSF, such as infection or neutropenia.

28. The use of a gene or DNA fragment according to any one of claims 1 to 8, a protein according to any one of claims 9 to 14, an antibody or its fragment according to any one of claims 15 to 17, a receptor according to claim 20 or a substance according to claim 22 or 23, in the production of a drug for diagnosis, prevention or treatment of a disease or condition related to G-CSF, such as

09937905 100101

infection or neutropenia.



ABSTRACT

The present invention provides, as a gene encoding an antigen recognized by G-CSF-inducing antibodies, a gene  
5 encoding:

(a) a protein having the amino acid sequence listed as SEQ. ID. No.2 of the Sequence Listing;

(b) a protein having the amino acid sequence listed as SEQ. ID. No.2 of the Sequence Listing with one or more  
10 amino acid deletions, substitutions, additions or insertions and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor; or

(c) a protein having at least 50% homology with the  
15 amino acid sequence listed as SEQ. ID. No.2 and also binding to an antibody or its fragment that is active to induce granulocyte colony-stimulating factor.

## SEQUENCE LISTING

&lt;110&gt; JAPAN TOBACCO INC.

&lt;120&gt; NOVEL PROTEINS, GENE ENCODING THE SAME AND METHOD OF UTILIZATION THEREOF

&lt;130&gt; YCT-486

&lt;160&gt; 4

&lt;210&gt; 1

&lt;211&gt; 840

&lt;212&gt; DNA

&lt;213&gt; Mouse macrophage cell RAW 264.7

&lt;400&gt;

gaacc atg tct ggc tgt caa gct caa gga gac tgt tgc tcg cgg ccg tgt 50

Met Ser Gly Cys Gln Ala Gln Gly Asp Cys Cys Ser Arg Pro Cys

1 5 10 15

ggc gcg cag gac aag gag cac ccc cga ttc ctg atc cca gaa ctt tgc 98

Gly Ala Gln Asp Lys Glu His Pro Arg Phe Leu Ile Pro Glu Leu Cys

20 25 30

aaa cag ttt tac cat ctg ggc tgg gtc act ggc act gga ggg gga atc 146

Lys Gln Phe Tyr His Leu Gly Trp Val Thr Gly Thr Gly Gly Gly Ile

35 40 45

agc ttg aag cat ggc aat gaa atc tac att gct ccc tca ggc gtg caa 194

Ser Leu Lys His Gly Asn Glu Ile Tyr Ile Ala Pro Ser Gly Val Gln

50 55 60

aag gag cgc att cag cca gaa gac atg ttt gtg tgt gac att aat gag 242

Lys Glu Arg Ile Gln Pro Glu Asp Met Phe Val Cys Asp Ile Asn Glu

65 70 75

cag gac ata agc ggg cct cca gca tct aag aag ctg aaa aaa agc cag 290

Gln Asp Ile Ser Gly Pro Pro Ala Ser Lys Lys Leu Lys Lys Ser Gln

80 85 90 95

tgc act cct ctt ttc atg aat gct tat acc atg aga gga gct ggc gca 338

Cys Thr Pro Leu Phe Met Asn Ala Tyr Thr Met Arg Gly Ala Gly Ala

100 105 110

gtg att cat acc cac tct aaa gct gct gtg atg gct acc ctt ctg ttt 386

Val Ile His Thr His Ser Lys Ala Ala Val Met Ala Thr Leu Leu Phe

115 120 125

$\begin{matrix} 1.1077 & 1.1225 & 1.1111 & 1.1225 & 1.1077 \\ 1.1077 & 1.1225 & 1.1111 & 1.1225 & 1.1077 \end{matrix}$

 $\langle 210 \rangle$  2

241

 $\langle 212 \rangle$  PRT $\langle 400 \rangle$ 

Met Ser Gly Cys Gln Ala Gln Gly Asp Cys Cys Ser Arg Pro Cys

1                      5                      10                      15

Gly Ala Gln Asp Lys Glu His Pro Arg Phe Leu Ile Pro Glu Leu Cys

20 25 30

Lys Gln Phe Tyr His Leu Gly Trp Val Thr Gly Thr Gly Gly Gly Ile

35                      40                      45



ggc gcg cag gac aag gag cat cca aga tac ctg atc cca gaa ctt tgc	141
Gly Ala Gln Asp Lys Glu His Pro Arg Tyr Leu Ile Pro Glu Leu Cys	
20 25 30	
aaa cag ttt tac cat tta ggc tgg gtc act ggg act gga gga gga att	189
Lys Gln Phe Tyr His Leu Gly Trp Val Thr Gly Thr Gly Gly Gly Ile	
35 40 45	
agc ttg aag cat ggc gat gaa atc tac att gct cct tca gga gtg caa	237
Ser Leu Lys His Gly Asp Glu Ile Tyr Ile Ala Pro Ser Gly Val Gln	
50 55 60	
aag gaa cga att cag cct gaa gac atg ttt gtt tat gat ata aat gaa	285
Lys Glu Arg Ile Gln Pro Glu Asp Met Phe Val Tyr Asp Ile Asn Glu	
65 70 75 80	
aag gac ata agt gga cct tgc cca tgc aag aag cta aaa aaa agc cag	333
Lys Asp Ile Ser Gly Pro Ser Pro Ser Lys Lys Leu Lys Lys Ser Gln	
85 90 95	
tgt act cct ctt ttc atg aat gct tac aca atg aga gga gca ggt gca	381
Cys Thr Pro Leu Phe Met Asn Ala Tyr Thr Met Arg Gly Ala Gly Ala	
100 105 110	
gtg att cat acc cac tct aaa gct gct gtg atg gcc acc ctt ctc ttt	429
Val Ile His Thr His Ser Lys Ala Ala Val Met Ala Thr Leu Leu Phe	
115 120 125	
cca gga cgg gag ttt aaa att aca cat caa gag atg ata aaa gga ata	477
Pro Gly Arg Glu Phe Lys Ile Thr His Gln Glu Met Ile Lys Gly Ile	
130 135 140	
aag aaa tgt act tcc gga ggg tat tat aga tat gat gat atg tta gtg	525
Lys Lys Cys Thr Ser Gly Gly Tyr Tyr Arg Tyr Asp Asp Met Leu Val	
145 150 155 160	
gta ccc att att gag aat aca cct gag gag aaa gac ctc aaa gat aga	573
Val Pro Ile Ile Glu Asn Thr Pro Glu Glu Lys Asp Leu Lys Asp Arg	
165 170 175	
atg gct cat gca atg aat gaa tac cca gac tcc tgt gca gta ctg gtc	621
Met Ala His Ala Met Asn Glu Tyr Pro Asp Ser Cys Ala Val Leu Val	
180 185 190	
aga cgt cat gga gta tat gtg tgg ggg gaa aca tgg gag aag gcc aaa	669
Arg Arg His Gly Val Tyr Val Trp Gly Glu Thr Trp Glu Lys Ala Lys	
195 200 205	

acc atg tgt gag tgt tat gac tat tta ttt gat att gcc gta tca atg	717
Thr Met Cys Glu Cys Tyr Asp Tyr Leu Phe Asp Ile Ala Val Ser Met	
210 215 220	
aag aaa gta gga ctt gat cct tca cag ctg cca gtt gga gaa aat gga	765
Lys Lys Val Gly Leu Asp Pro Ser Gln Leu Pro Val Gly Glu Asn Gly	
225 230 235 240	
att gtc taa gccaaaagaa agtctaatta tatacagaga taaagctaaa	814
Ile Val	
cgtaattatt atttaaatga aagctatttt tttaaatgaa ttgaaatttt tcatgatgct	874
actaatttgc cactaaatac tgcaaatggg caccctgaat ctcttcctgac	924

 $\langle 210 \rangle$  4

&lt;211&gt; 242

&lt;212&gt; PRT

 $\langle 400 \rangle$ 

Met	Ser	Gly	Cys	Asp	Ala	Trp	Glu	Gly	Asp	Cys	Cys	Ser	Arg	Arg	Cys
1				5					10					15	
Gly	Ala	Gln	Asp	Lys	Glu	His	Pro	Arg	Tyr	Leu	Ile	Pro	Glu	Leu	Cys
			20					25					30		
Lys	Gln	Phe	Tyr	His	Leu	Gly	Trp	Val	Thr	Gly	Thr	Gly	Gly	Gly	Ile
		35					40					45			
Ser	Leu	Lys	His	Gly	Asp	Glu	Ile	Tyr	Ile	Ala	Pro	Ser	Gly	Val	Gln
	50					55				60					
Lys	Glu	Arg	Ile	Gln	Pro	Glu	Asp	Met	Phe	Val	Tyr	Asp	Ile	Asn	Glu
65				70					75					80	
Lys	Asp	Ile	Ser	Gly	Pro	Ser	Pro	Ser	Lys	Lys	Leu	Lys	Lys	Ser	Gln
			85					90					95		
Cys	Thr	Pro	Leu	Phe	Met	Asn	Ala	Tyr	Thr	Met	Arg	Gly	Ala	Gly	Ala
		100						105				110			
Val	Ile	His	Thr	His	Ser	Lys	Ala	Ala	Val	Met	Ala	Thr	Leu	Leu	Phe
	115					120					125				
Pro	Gly	Arg	Glu	Phe	Lys	Ile	Thr	His	Gln	Glu	Met	Ile	Lys	Gly	Ile
	130					135				140					
Lys	Lys	Cys	Thr	Ser	Gly	Gly	Tyr	Tyr	Arg	Tyr	Asp	Asp	Met	Leu	Val
145				150					155					160	
Val	Pro	Ile	Ile	Glu	Asn	Thr	Pro	Glu	Glu	Lys	Asp	Leu	Lys	Asp	Arg

	165		170		175	
Met	Ala	His	Ala	Met	Asn	Glu Tyr Pro Asp Ser Cys Ala Val Leu Val
	180		185		190	
Arg	Arg	His	Gly	Val	Tyr	Val Trp Gly Glu Thr Trp Glu Lys Ala Lys
	195		200		205	
Thr	Met	Cys	Glu	Cys	Tyr	Asp Tyr Leu Phe Asp Ile Ala Val Ser Met
	210		215		220	
Lys	Lys	Val	Gly	Leu	Asp	Pro Ser Gln Leu Pro Val Gly Glu Asn Gly
225			230		235	240
Ile	Val					



ENTERED

PCT

## RAW SEQUENCE LISTING

PATENT APPLICATION: US/09/937,905

DATE: 12/02/2002

TIME: 15:58:08

Input Set : A:\0230-0169P.st25.txt

Output Set: N:\CRF4\12022002\I937905.raw

3 <110> APPLICANT: SHA, Shiken  
 4 AOKI, Yoshiko  
 5 NISHI, Yoshisuke  
 7 <120> TITLE OF INVENTION: Novel Proteins, Gene Encoding the Same and Method of Utilization Thereof  
 9 <130> FILE REFERENCE: 0230-0169P  
 11 <140> CURRENT APPLICATION NUMBER: US 09/937,905  
 C--> 12 <141> CURRENT FILING DATE: 2002-01-14  
 14 <150> PRIOR APPLICATION NUMBER: PCT/JP00/02080  
 15 <151> PRIOR FILING DATE: 2000-03-31  
 17 <150> PRIOR APPLICATION NUMBER: Japan 95092/1999  
 18 <151> PRIOR FILING DATE: 1999-04-01  
 20 <160> NUMBER OF SEQ ID NOS: 8  
 22 <170> SOFTWARE: PatentIn version 3.1  
 24 <210> SEQ ID NO: 1  
 25 <211> LENGTH: 840  
 26 <212> TYPE: DNA  
 27 <213> ORGANISM: Mus sp.  
 29 <220> FEATURE:  
 30 <221> NAME/KEY: CDS  
 31 <222> LOCATION: (6)..(731)  
 32 <223> OTHER INFORMATION:  
 35 <220> FEATURE:  
 36 <221> NAME/KEY: misc\_feature  
 37 <222> LOCATION: (1)..(840)  
 38 <223> OTHER INFORMATION: Mouse macrophage cell RAW 264.7  
 W--> 41 <400> 1  
 42 gaacc atg tct ggc tgt caa gct caa gga gac tgt tgc tcg cgg ccg tgt 50  
 43 Met Ser Gly Cys Gln Ala Gln Gly Asp Cys Cys Ser Arg Pro Cys  
 44 1 5 10 15  
 46 ggc gcg cag gac aag gag cac ccc cga ttc ctg atc cca gaa ctt tgc 98  
 47 Gly Ala Gln Asp Lys Glu His Pro Arg Phe Leu Ile Pro Glu Leu Cys  
 48 20 25 30  
 50 aaa cag ttt tac cat ctg ggc tgg gtc act ggc act gga ggg gga atc 146  
 51 Lys Gln Phe Tyr His Leu Gly Trp Val Thr Gly Thr Gly Gly Gly Ile  
 52 35 40 45  
 54 agc ttg aag cat ggc aat gaa atc tac att gct ccc tca ggc gtg caa 194  
 55 Ser Leu Lys His Gly Asn Glu Ile Tyr Ile Ala Pro Ser Gly Val Gln  
 56 50 55 60  
 58 aag gag cgc att cag cca gaa gac atg ttt gtg tgt gac att aat gag 242  
 59 Lys Glu Arg Ile Gln Pro Glu Asp Met Phe Val Cys Asp Ile Asn Glu  
 60 65 70 75  
 62 cag gac ata agc ggg cct cca gca tct aag aag ctg aaa aaa agc cag 290  
 63 Gln Asp Ile Ser Gly Pro Pro Ala Ser Lys Lys Leu Lys Lys Ser Gln



## RAW SEQUENCE LISTING

PATENT APPLICATION: US/09/937,905

DATE: 12/02/2002

TIME: 15:58:08

Input Set : A:\0230-0169P.st25.txt

Output Set: N:\CRF4\12022002\I937905.raw

```

64 80          85          90          95
66 tgc act cct ctt ttc atg aat gct tat acc atg aga gga gct ggc gca      338
67 Cys Thr Pro Leu Phe Met Asn Ala Tyr Thr Met Arg Gly Ala Gly Ala
68          100          105          110
70 gtg att cat acc cac tct aaa gct gct gtg atg gct acc ctt ctg ttt      386
71 Val Ile His Thr His Ser Lys Ala Ala Val Met Ala Thr Leu Leu Phe
72          115          120          125
74 cca gga cag gag ttt aaa att aca cat caa gag atg atc aaa gga ata      434
75 Pro Gly Gln Glu Phe Lys Ile Thr His Gln Glu Met Ile Lys Gly Ile
76          130          135          140
78 agg aaa tgt acc tca gga ggc tat tac aga tac gat gat atg tta gtg      482
79 Arg Lys Cys Thr Ser Gly Gly Tyr Tyr Arg Tyr Asp Asp Met Leu Val
80          145          150          155
82 gta cct att att gag aac act cct gaa gag aag gat ctc aaa gaa agg      530
83 Val Pro Ile Ile Glu Asn Thr Pro Glu Glu Lys Asp Leu Lys Glu Arg
84 160          165          170          175
86 atg gct cat gcc atg aat gag tac cca gac tcc tgt gcg gtt ctt gtc      578
87 Met Ala His Ala Met Asn Glu Tyr Pro Asp Ser Cys Ala Val Leu Val
88          180          185          190
90 cgg cgt cat ggg gtg tac gtg tgg gga gaa aca tgg gag aaa gca aaa      626
91 Arg Arg His Gly Val Tyr Val Trp Gly Glu Thr Trp Glu Lys Ala Lys
92          195          200          205
94 acc atg tgt gag tgt tat gac tac ctg ttt gac att gct gtc tcc atg      674
95 Thr Met Cys Glu Cys Tyr Asp Tyr Leu Phe Asp Ile Ala Val Ser Met
96          210          215          220
98 aag aag atg gga ctc gat cca aca cag ctc cca gtt gga gaa aat gga      722
99 Lys Lys Met Gly Leu Asp Pro Thr Gln Leu Pro Val Gly Glu Asn Gly
100          225          230          235
102 att gtg taa gccaaagtga tgcctaagca tctccaacaa taaaacaaac      771
103 Ile Val
104 240
106 tcaattatgc cttaaataaaa actcagctgc ttttaaaaaa aaaaaaaaaa aaaaaaaaaa      831
108 aaaaaaaaaa      840
111 <210> SEQ ID NO: 2
112 <211> LENGTH: 241
113 <212> TYPE: PRT
114 <213> ORGANISM: Mus sp.
116 <220> FEATURE:
117 <221> NAME/KEY: misc_feature
118 <222> LOCATION: (1)..(241)
119 <223> OTHER INFORMATION: Mouse macrophage cell RAW 264.7
121 <400> SEQUENCE: 2
123 Met Ser Gly Cys Gln Ala Gln Gly Asp Cys Cys Ser Arg Pro Cys Gly
124 1          5          10          15
127 Ala Gln Asp Lys Glu His Pro Arg Phe Leu Ile Pro Glu Leu Cys Lys
128          20          25          30
131 Gln Phe Tyr His Leu Gly Trp Val Thr Gly Thr Gly Gly Gly Ile Ser
132          35          40          45
135 Leu Lys His Gly Asn Glu Ile Tyr Ile Ala Pro Ser Gly Val Gln Lys

```

## RAW SEQUENCE LISTING

PATENT APPLICATION: US/09/937,905

DATE: 12/02/2002

TIME: 15:58:08

Input Set : A:\0230-0169P.st25.txt

Output Set: N:\CRF4\12022002\I937905.raw

```

136      50      55      60
139 Glu Arg Ile Gln Pro Glu Asp Met Phe Val Cys Asp Ile Asn Glu Gln
140 65      70      75      80
143 Asp Ile Ser Gly Pro Pro Ala Ser Lys Lys Leu Lys Lys Ser Gln Cys
144      85      90      95
147 Thr Pro Leu Phe Met Asn Ala Tyr Thr Met Arg Gly Ala Gly Ala Val
148      100      105      110
151 Ile His Thr His Ser Lys Ala Ala Val Met Ala Thr Leu Leu Phe Pro
152      115      120      125
155 Gly Gln Glu Phe Lys Ile Thr His Gln Glu Met Ile Lys Gly Ile Arg
156      130      135      140
159 Lys Cys Thr Ser Gly Gly Tyr Tyr Arg Tyr Asp Asp Met Leu Val Val
160 145      150      155      160
163 Pro Ile Ile Glu Asn Thr Pro Glu Glu Lys Asp Leu Lys Glu Arg Met
164      165      170      175
167 Ala His Ala Met Asn Glu Tyr Pro Asp Ser Cys Ala Val Leu Val Arg
168      180      185      190
171 Arg His Gly Val Tyr Val Trp Gly Glu Thr Trp Glu Lys Ala Lys Thr
172      195      200      205
175 Met Cys Glu Cys Tyr Asp Tyr Leu Phe Asp Ile Ala Val Ser Met Lys
176      210      215      220
179 Lys Met Gly Leu Asp Pro Thr Gln Leu Pro Val Gly Glu Asn Gly Ile
180 225      230      235      240
183 Val

```

187 &lt;210&gt; SEQ ID NO: 3

188 &lt;211&gt; LENGTH: 924

189 &lt;212&gt; TYPE: DNA

190 &lt;213&gt; ORGANISM: Homo sapiens

192 &lt;220&gt; FEATURE:

193 &lt;221&gt; NAME/KEY: CDS

194 &lt;222&gt; LOCATION: (46)..(774)

195 &lt;223&gt; OTHER INFORMATION:

198 &lt;220&gt; FEATURE:

199 &lt;221&gt; NAME/KEY: misc\_feature

200 &lt;222&gt; LOCATION: (1)..(924)

201 &lt;223&gt; OTHER INFORMATION: Human normal brain tissue

W--&gt; 204 &lt;400&gt; 3

```

205 aagccgtgcg gagattggag gccgcgcggg tccctggtct gggcc atg tct ggc.tgt      57
206                               Met Ser Gly Cys
207                               1
209 gat gct tgg gag gga gac tgt tgt tcc cgg aga tgc ggc gcg cag gac      105
210 Asp Ala Trp Glu Gly Asp Cys Cys Ser Arg Arg Cys Gly Ala Gln Asp
211 5      10      15      20
213 aag gag cat cca aga tac ctg atc cca gaa ctt tgc aaa cag ttt tac      153
214 Lys Glu His Pro Arg Tyr Leu Ile Pro Glu Leu Cys Lys Gln Phe Tyr
215      25      30      35
217 cat tta ggc tgg gtc act ggg act gga gga gga att agc ttg aag cat      201
218 His Leu Gly Trp Val Thr Gly Thr Gly Gly Gly Ile Ser Leu Lys His
219      40      45      50

```

## RAW SEQUENCE LISTING

PATENT APPLICATION: US/09/937,905

DATE: 12/02/2002

TIME: 15:58:08

Input Set : A:\0230-0169P.st25.txt

Output Set: N:\CRF4\12022002\I937905.raw

```

221 ggc gat gaa atc tac att gct cct tca gga gtg caa aag gaa cga att      249
222 Gly Asp Glu Ile Tyr Ile Ala Pro Ser Gly Val Gln Lys Glu Arg Ile
223      55      60      65
225 cag cct gaa gac atg ttt gtt tat gat ata aat gaa aag gac ata agt      297
226 Gln Pro Glu Asp Met Phe Val Tyr Asp Ile Asn Glu Lys Asp Ile Ser
227      70      75      80
229 gga cct tcg cca tcg aag aag cta aaa aaa agc cag tgt act cct ctt      345
230 Gly Pro Ser Pro Ser Lys Lys Leu Lys Lys Ser Gln Cys Thr Pro Leu
231 85      90      95      100
233 ttc atg aat gct tac aca atg aga gga gca ggt gca gtg att cat acc      393
234 Phe Met Asn Ala Tyr Thr Met Arg Gly Ala Gly Ala Val Ile His Thr
235      105      110      115
237 cac tct aaa gct gct gtg atg gcc acc ctt ctc ttt cca gga cgg gag      441
238 His Ser Lys Ala Ala Val Met Ala Thr Leu Leu Phe Pro Gly Arg Glu
239      120      125      130
241 ttt aaa att aca cat caa gag atg ata aaa gga ata aag aaa tgt act      489
242 Phe Lys Ile Thr His Gln Glu Met Ile Lys Gly Ile Lys Lys Cys Thr
243      135      140      145
245 tcc gga ggg tat tat aga tat gat gat atg tta gtg gta ccc att att      537
246 Ser Gly Gly Tyr Tyr Arg Tyr Asp Asp Met Leu Val Val Pro Ile Ile
247      150      155      160
249 gag aat aca cct gag gag aaa gac ctc aaa gat aga atg gct cat gca      585
250 Glu Asn Thr Pro Glu Glu Lys Asp Leu Lys Asp Arg Met Ala His Ala
251 165      170      175      180
253 atg aat gaa tac cca gac tcc tgt gca gta ctg gtc aga cgt cat gga      633
254 Met Asn Glu Tyr Pro Asp Ser Cys Ala Val Leu Val Arg Arg His Gly
255      185      190      195
257 gta tat gtg tgg ggg gaa aca tgg gag aag gcc aaa acc atg tgt gag      681
258 Val Tyr Val Trp Gly Glu Thr Trp Glu Lys Ala Lys Thr Met Cys Glu
259      200      205      210
261 tgt tat gac tat tta ttt gat att gcc gta tca atg aag aaa gta gga      729
262 Cys Tyr Asp Tyr Leu Phe Asp Ile Ala Val Ser Met Lys Lys Val Gly
263      215      220      225
265 ctt gat cct tca cag ctc cca gtt gga gaa aat gga att gtc taa      774
266 Leu Asp Pro Ser Gln Leu Pro Val Gly Glu Asn Gly Ile Val
267      230      235      240
269 gccaaaagaa agtctaatta tatacagaga taaagctaaa cgtaattatt atttaaata      834
271 aagctat tttttaaataa ttgaaat tcatgatgct actaatttgc cactaaatac      894
273 tgcaaaggt caccctgaat ctcttctgac      924
276 <210> SEQ ID NO: 4
277 <211> LENGTH: 242
278 <212> TYPE: PRT
279 <213> ORGANISM: Homo sapiens
281 <220> FEATURE:
282 <221> NAME/KEY: misc_feature
283 <222> LOCATION: (1)..(242)
284 <223> OTHER INFORMATION: Human normal brain tissue
286 <400> SEQUENCE: 4
288 Met Ser Gly Cys Asp Ala Trp Glu Gly Asp Cys Cys Ser Arg Arg Cys

```

## RAW SEQUENCE LISTING

PATENT APPLICATION: US/09/937,905

DATE: 12/02/2002

TIME: 15:58:08

Input Set : A:\0230-0169P.st25.txt

Output Set: N:\CRF4\12022002\I937905.raw

```

289 1          5          10          15
292 Gly Ala Gln Asp Lys Glu His Pro Arg Tyr Leu Ile Pro Glu Leu Cys
293          20          25          30
296 Lys Gln Phe Tyr His Leu Gly Trp Val Thr Gly Thr Gly Gly Gly Ile
297          35          40          45
300 Ser Leu Lys His Gly Asp Glu Ile Tyr Ile Ala Pro Ser Gly Val Gln
301          50          55          60
304 Lys Glu Arg Ile Gln Pro Glu Asp Met Phe Val Tyr Asp Ile Asn Glu
305 65          70          75          80
308 Lys Asp Ile Ser Gly Pro Ser Pro Ser Lys Lys Leu Lys Lys Ser Gln
309          85          90          95
312 Cys Thr Pro Leu Phe Met Asn Ala Tyr Thr Met Arg Gly Ala Gly Ala
313          100          105          110
316 Val Ile His Thr His Ser Lys Ala Ala Val Met Ala Thr Leu Leu Phe
317          115          120          125
320 Pro Gly Arg Glu Phe Lys Ile Thr His Gln Glu Met Ile Lys Gly Ile
321          130          135          140
324 Lys Lys Cys Thr Ser Gly Gly Tyr Tyr Arg Tyr Asp Asp Met Leu Val
325 145          150          155          160
328 Val Pro Ile Ile Glu Asn Thr Pro Glu Glu Lys Asp Leu Lys Asp Arg
329          165          170          175
332 Met Ala His Ala Met Asn Glu Tyr Pro Asp Ser Cys Ala Val Leu Val
333          180          185          190
336 Arg Arg His Gly Val Tyr Val Trp Gly Glu Thr Trp Glu Lys Ala Lys
337          195          200          205
340 Thr Met Cys Glu Cys Tyr Asp Tyr Leu Phe Asp Ile Ala Val Ser Met
341          210          215          220
344 Lys Lys Val Gly Leu Asp Pro Ser Gln Leu Pro Val Gly Glu Asn Gly
345 225          230          235          240
348 Ile Val

```

352 &lt;210&gt; SEQ ID NO: 5

353 &lt;211&gt; LENGTH: 19

354 &lt;212&gt; TYPE: DNA

355 &lt;213&gt; ORGANISM: Artificial Sequence

357 &lt;220&gt; FEATURE:

358 &lt;223&gt; OTHER INFORMATION: Sense primer of position 4 to 22 of the mouse antigen gene

MMR19

360 &lt;400&gt; SEQUENCE: 5

361 ccatgtcttg ctgtcaagc

19

364 &lt;210&gt; SEQ ID NO: 6

365 &lt;211&gt; LENGTH: 21

366 &lt;212&gt; TYPE: DNA

367 &lt;213&gt; ORGANISM: Artificial Sequence

369 &lt;220&gt; FEATURE:

370 <223> OTHER INFORMATION: An antisense primer of position 714 to 724 of the mouse  
anitgen g

371 ene MMR19

373 &lt;400&gt; SEQUENCE: 6

374 ccattttctc caactgggag c

21

377 &lt;210&gt; SEQ ID NO: 7

378 &lt;211&gt; LENGTH: 23

RAW SEQUENCE LISTING ERROR SUMMARY  
PATENT APPLICATION: US/09/937,905

DATE: 12/02/2002  
TIME: 15:58:09

Input Set : A:\0230-0169P.st25.txt  
Output Set: N:\CRF4\12022002\I937905.raw

Invalid Line Length:

The rules require that a line not exceed 72 characters in length. This includes spaces.

Seq#:1; Line(s) 7

**VERIFICATION SUMMARY**

PATENT APPLICATION: US/09/937,905

DATE: 12/02/2002

TIME: 15:58:09

Input Set : A:\0230-0169P.st25.txt

Output Set: N:\CRF4\12022002\I937905.raw

L:12 M:271 C: Current Filing Date differs, Replaced Current Filing Date  
L:41 M:258 W: Mandatory Feature missing, <223> Blank for SEQ#:1,Line#:32  
L:204 M:258 W: Mandatory Feature missing, <223> Blank for SEQ#:3,Line#:195

**BIRCH, STEWART, KOŁASCH & BIRCH, LLP****COMBINED DECLARATION AND POWER OF ATTORNEY**

ATTORNEY DOCKET NO.

0230-0169P

PLEASE NOTE:  
YOU MUST  
COMPLETE THE  
FOLLOWING:

Insert Title:

NOVEL PROTEINS, GENE ENCODING THE SAME AND METHOD OF UTILIZATION  
THEREOF

Fill in Appropriate  
Information -  
For Use Without  
Specification  
Attached:

the specification of which is attached hereto. If not attached hereto,

the specification was filed on \_\_\_\_\_ as  
United States Application Number \_\_\_\_\_; and /or

the specification was filed on March 31, 2000 as PCT  
International Application Number PCT/JP00/02080; and was  
amended under PCT Article 19 on \_\_\_\_\_ (if applicable)

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations, §1.56.

I do not know and do not believe the same was ever known or used in the United States of America before my or our invention thereof, or patented or described in any printed publication in any country before my or our invention thereof or more than one year prior to this application, that the same was not in public use or on sale in the United States of America more than one year prior to this application, that the invention has not been patented or made the subject of an inventor's certificate issued before the date of this application in any country foreign to the United States of America on an application filed by me or my legal representatives or assigns more than twelve months (six months for designs) prior to this application, and that no application for patent or inventor's certificate on this invention has been filed in any country foreign to the United States of America prior to this application by me or my legal representatives or assigns, except as follows.

I hereby claim foreign priority benefits under Title 35, United States Code, §119 (a)-(d) of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed:

Insert Priority  
Information:  
(if appropriate)

**Prior Foreign Application(s)**

<u>95092/1999</u> (Number)	<u>Japan</u> (Country)	<u>4/1/1999</u> (Month/Day/Year Filed)
_____ (Number)	_____ (Country)	_____ (Month/Day/Year Filed)
_____ (Number)	_____ (Country)	_____ (Month/Day/Year Filed)
_____ (Number)	_____ (Country)	_____ (Month/Day/Year Filed)
_____ (Number)	_____ (Country)	_____ (Month/Day/Year Filed)

**Priority Claimed**

<input checked="" type="checkbox"/> Yes	<input type="checkbox"/> No
<input type="checkbox"/> Yes	<input type="checkbox"/> No
<input type="checkbox"/> Yes	<input type="checkbox"/> No
<input type="checkbox"/> Yes	<input type="checkbox"/> No
<input type="checkbox"/> Yes	<input type="checkbox"/> No

Insert Provisional  
Application(s):  
(if any)

I hereby claim the benefit under Title 35, United States Code, §119(e) of any United States provisional application(s) listed below.

_____ (Application Number)	_____ (Filing Date)
_____ (Application Number)	_____ (Filing Date)

All Foreign Applications, if any, for any Patent or Inventor's Certificate Filed More Than 12 Months (6 Months for Designs) Prior To The Filing Date of This Application:

Insert Requested  
Information:  
(if appropriate)

Country	Application No.	Date of Filing (Month/Day/Year)
_____	_____	_____
_____	_____	_____

I hereby claim the benefit under Title 35, United States Code, §120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations, §1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application:

Insert Prior U.S.  
Application(s):  
(if any)

_____ (Application Number)	_____ (Filing Date)	_____ (Status - patented, pending, abandoned)
_____ (Application Number)	_____ (Filing Date)	_____ (Status - patented, pending, abandoned)

I hereby appoint the following attorneys to prosecute this application and/or an international application based on this application and to transact all business in the Patent and Trademark Office connected therewith and in connection with the resulting patent based on instructions received from the entity who first sent the application papers to the attorneys identified below, unless the inventor(s) or assignee provides said attorneys with a written notice to the contrary:

Terrell C. Birch (Reg. No. 19,382)  
 Joseph A. Kolasch (Reg. No. 22,463)  
 Bernard L. Sweeney (Reg. No. 24,448)  
 Charles Gorenstein (Reg. No. 29,271)  
 Leonard R. Svensson (Reg. No. 30,330)  
 Andrew D. Meikle (Reg. No. 32,868)  
 Joe McKinney Muncy (Reg. No. 32,334)  
 C. Joseph Faraci (Reg. No. 32,350)

Raymond C. Stewart (Reg. No. 21,066)  
 James M. Slattery (Reg. No. 28,380)  
 Michael K. Mutter (Reg. No. 29,680)  
 Gerald M. Murphy, Jr. (Reg. No. 28,977)  
 Terry L. Clark (Reg. No. 32,644)  
 Marc S. Weiner (Reg. No. 32,181)  
 Andrew F. Reish (Reg. No. 33,443)  
 Donald J. Daley (Reg. No. 34,313)

Send Correspondence to:

**BIRCH, STEWART, KOLASCH & BIRCH, LLP**

P.O. Box 747 • Falls Church, Virginia 22040-0747

Telephone: (703) 205-8000 • Facsimile: (703) 205-8050

PLEASE NOTE:  
 YOU MUST  
 COMPLETE THE  
 FOLLOWING:

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Full Name of First or Sole  
 Inventor:  
 Insert Name of Inventor  
 Insert Date This  
 Document is Signed

Insert Residence  
 Insert Citizenship

Insert Post Office  
 Address

Full Name of Second  
 Inventor, if any:

see above

Full Name of Third  
 Inventor, if any

see above

Full Name of Fourth  
 Inventor, if any

see above

Full Name of Fifth  
 Inventor, if any

see above

GIVEN NAME	FAMILY NAME	INVENTOR'S SIGNATURE	DATE*
Shiken	SHA	Shiken Sha	Sep. 21, 2001
Residence (City, State & Country)		CITIZENSHIP	
Kanagawa, Japan		Chinese	
POST OFFICE ADDRESS (Complete Street Address including City, State & Country)			
c/o Affiliated Business Division Aobadai, Japan Tobacco Inc. of 6-2, Umegaoka, Aoba-ku, Yokohama-shi, Kanagawa 227-8512 Japan SPK			
GIVEN NAME	FAMILY NAME	INVENTOR'S SIGNATURE	DATE*
Yoshiko	AOKI	Yoshiko Aoki	Sep 21, 2001
Residence (City, State & Country)		CITIZENSHIP	
Kanagawa, Japan		Japanese	
POST OFFICE ADDRESS (Complete Street Address including City, State & Country)			
c/o Affiliated Business Division Aobadai, Japan Tobacco Inc. of 6-2, Umegaoka, Aoba-ku, Yokohama-shi, Kanagawa 227-8512 Japan SPK			
GIVEN NAME	FAMILY NAME	INVENTOR'S SIGNATURE	DATE*
Yoshisuke	NISHI	Yoshisuke Nishi	Sep. 21, 2001
Residence (City, State & Country)		CITIZENSHIP	
Kanagawa, Japan		Japanese	
POST OFFICE ADDRESS (Complete Street Address including City, State & Country)			
c/o Affiliated Business Division Aobadai, Japan Tobacco Inc. of 6-2, Umegaoka, Aoba-ku, Yokohama-shi, Kanagawa 227-8512 Japan SPK			
GIVEN NAME	FAMILY NAME	INVENTOR'S SIGNATURE	DATE*
Residence (City, State & Country)		CITIZENSHIP	
POST OFFICE ADDRESS (Complete Street Address including City, State & Country)			
GIVEN NAME	FAMILY NAME	INVENTOR'S SIGNATURE	DATE*
Residence (City, State & Country)		CITIZENSHIP	
POST OFFICE ADDRESS (Complete Street Address including City, State & Country)			